

ON MOMENTS OF A POLYTOPE

NICK GRAVIN, DMITRII V. PASECHNIK, BORIS SHAPIRO, AND MICHAEL SHAPIRO

To the memory of Mikael Passare

ABSTRACT. We show that the multivariate generating function of appropriately normalized moments of a measure with homogeneous polynomial density supported on a compact polytope $\mathcal{P} \subset \mathbb{R}^d$ is a rational function. Its denominator is the product of linear forms dual to the vertices of $\mathcal{P}$ raised to the power equal to the degree of the density function. Using this, we solve the inverse moment problem for the set of, not necessarily convex, polytopes having a given set S of vertices. Under a weak non-degeneracy assumption we also show that the uniform measure supported on any such polytope is a linear combination of uniform measures supported on simplices with vertices in S .

1. INTRODUCTION

The initial motivation for the present paper came from proposed in [16] efficient algorithm recovering an arbitrary convex polytope from axial moments of a polynomial measure supported on it. This algorithm is based on the formulas for the axial moments of polytopes found over 20 years ago independently by M. Brion, J. Lawrence, A. Khovanskii-A. Pukhlikov, and A. Barvinok [6, 11, 18, 23], see [7, 8] for accessible explanation. In [16] the authors made an essential, although implicit, use of a univariate rational generating function for appropriately normalized axial moments. Here a multivariate, and explicit, analog of the latter function is developed. It turns out it provides a very convenient encoding of non-convex polytopes, which is of independent interest. E.g. it leads to a natural definition of *vertices* of such non-convex polytopes, which have similar properties to vertices of convex polytopes. It also allows to find the *exact solutions* of a class of inverse moment problems on non-convex polytopes.

Notation. In what follows we shall always assume that $\mathbb{R}^d$ is endowed with a fixed coordinate system $(x_1, \dots, x_d)$, orthonormal with respect to the standard scalar product $\langle \cdot, \cdot \rangle$. Let μ be a finite complex-valued Borel measure in $\mathbb{R}^d$. (For standard measure-theoretic notions we follow [25].) Given a multiindex $I = (i_1, \dots, i_d)$, let $\mathbf{x}^I$ be the shorthand of the monomial $x_1^{i_1} \dots x_d^{i_d}$ and $|I|$ the shorthand for $i_1 + \dots + i_d$. For any multiindex I , define the *moment* $m_I(\mu)$ of μ as

$$m_I(\mu) := \int_{\mathbb{R}^d} x_1^{i_1} x_2^{i_2} \dots x_d^{i_d} d\mu(x_1, x_2, \dots, x_d) = \int_{\mathbb{R}^d} \mathbf{x}^I d\mu(\mathbf{x}). \quad (1.1)$$

Define the *normalized moment generating function* $F_\mu(\mathbf{u}) = F_\mu(u_1, \dots, u_d)$ of μ by

$$F_\mu(\mathbf{u}) := \sum_{I=(i_1, \dots, i_d) \geq 0} \frac{(|I| + d)!}{i_1! \dots i_d!} m_I(\mu) \mathbf{u}^I, \quad \text{where } \mathbf{u}^I = u_1^{i_1} \dots u_d^{i_d}. \quad (1.2)$$

Date: October 12, 2012.

2010 Mathematics Subject Classification. Primary 44A60; Secondary 31B20.

Key words and phrases. moments of a polytope, generating function.

Note that $F_\mu(\mathbf{u})$ admits the integral representation

$$F_\mu(\mathbf{u}) = d! \int_{\mathbb{R}^d} \frac{d\mu(\mathbf{x})}{(1 - \langle \mathbf{x}, \mathbf{u} \rangle)^{d+1}}, \quad (1.3)$$

which is a special case of a *Fantappiè transformation*. For details on the latter, see e.g. [4, Chapter 3]. A proof of (1.3) will be given at the end of Section 2; see also Remark 10.

Given any complex-valued finite measure μ and any degree δ homogeneous d -variate polynomial ρ , it is convenient to define the (re)normalized moment generating function $F_\mu^\rho(\mathbf{u})$ for the measure $\rho\mu$, where by definition, $\int_{\mathbb{R}^d} f d(\rho\mu) = \int_{\mathbb{R}^d} f \rho d\mu$, in such a way that it can be obtained from $F_\mu(u)$ by application of the differential operator $\rho \left(\frac{\partial}{\partial \mathbf{u}} \right)$. Namely, set

$$F_\mu^\rho(\mathbf{u}) := \sum_{I:=(i_1, \dots, i_d) \geq 0} \frac{(|I| + d + \delta)!}{i_1! \cdots i_d!} m_I(\rho\mu) \mathbf{u}^I. \quad (1.4)$$

Note that $F_\mu^\rho(\mathbf{u}) \neq F_{\rho\mu}(\mathbf{u})$ for non-constant ρ . However, they are also connected, by an explicit differential operator as follows.

Theorem 1. For any complex-valued finite measure μ and any homogeneous polynomial ρ of degree δ ,

$$F_\mu^\rho(\mathbf{u}) = \prod_{\ell=d}^{d+\delta-1} \left(\sum_k u_k \frac{\partial}{\partial u_k} + \ell \right) \circ F_{\rho\mu}(\mathbf{u}) \quad (1.5)$$

$$= \rho \left(\frac{\partial}{\partial \mathbf{u}} \right) \circ F_\mu(\mathbf{u}) \quad (1.6)$$

$$= (d + \delta)! \int_{\mathbb{R}^d} \frac{\rho(\mathbf{x}) d\mu(\mathbf{x})}{(1 - \langle \mathbf{x}, \mathbf{u} \rangle)^{d+\delta+1}}. \quad (1.7)$$

Here and in what follows $\circ$ denotes the application of a differential operator to a function. The proof of the latter result is basically an exercise in manipulating formal power series, and we do not claim its novelty. For the sake of completeness, we include a proof in Section 2.

Results on convex polytopes. A finite set $S \subset \mathbb{R}^d$ is called *spanning* if it is not contained in any (affine) hyperplane in $\mathbb{R}^d$. (Obviously, $\text{card}(S) \geq d + 1$.) As usual, by a (compact, convex) *polytope* $\mathcal{P} \subset \mathbb{R}^d$ we mean the convex hull of a finite spanning set in $\mathbb{R}^d$. The set of vertices of a convex polytope $\mathcal{P}$ is the inclusion-minimal finite set with convex hull $\mathcal{P}$. A d -*simplex* in $\mathbb{R}^d$ is the convex hull of a spanning $(d + 1)$ -tuple of points. By an *open polytope* (resp. *simplex*) we mean the set of interior points of a compact polytope (resp. simplex).

Given a convex polytope $\mathcal{P}$ let $\mathcal{V} = (\mathbf{v}_1, \dots, \mathbf{v}_N)$ denote the set of its vertices. Assume that $\mathcal{P}$ is simple, i.e. each $\mathbf{v} \in \mathcal{V}$ has exactly d incident edges $\mathbf{v}\mathbf{v}_{e_1}, \dots, \mathbf{v}\mathbf{v}_{e_d}$. Set $w_k(\mathbf{v}) := \mathbf{v}_{e_k} - \mathbf{v}$, for $1 \leq k \leq d$. The non-negative real span $K_{\mathbf{v}}$ of $w_1(\mathbf{v}), \dots, w_d(\mathbf{v})$ is called the *tangent cone* of $\mathcal{P}$ at $\mathbf{v}$. For each $K_{\mathbf{v}}$ define $|\det K_{\mathbf{v}}| = |\det(w_1(\mathbf{v}), \dots, w_d(\mathbf{v}))|$ to be the volume of the parallelepiped formed by $w_1(\mathbf{v}), \dots, w_d(\mathbf{v})$.

Given a bounded domain $\Omega \subset \mathbb{R}^d$, we call the measure

$$\mu_\Omega = \chi_\Omega dx_1 dx_2 \dots dx_d,$$

where χ_Ω is the characteristic function of Ω , the *standard measure* of Ω .

For a simple convex polytope $\mathcal{P}$ we have the following explicit representation of $F_{\mu_{\mathcal{P}}}(\mathbf{u})$.

Theorem 2. For an arbitrary simple convex polytope $\mathcal{P}$,

$$F_{\mathcal{P}}(\mathbf{u}) := F_{\mu_{\mathcal{P}}}(\mathbf{u}) = (-1)^d \sum_{\mathbf{v} \in \mathcal{V}} \frac{\langle \mathbf{v}, \mathbf{u} \rangle^d |\det K_{\mathbf{v}}|}{\prod_{j=1}^d \langle w_j(\mathbf{v}), \mathbf{u} \rangle} \cdot \frac{1}{1 - \langle \mathbf{v}, \mathbf{u} \rangle} \quad (1.8)$$

$$= (-1)^d \sum_{\mathbf{v} \in \mathcal{V}} \frac{|\det K_{\mathbf{v}}|}{\prod_{j=1}^d \langle w_j(\mathbf{v}), \mathbf{u} \rangle} \cdot \frac{1}{1 - \langle \mathbf{v}, \mathbf{u} \rangle}. \quad (1.9)$$

Remark 1. Instead of the explicit choice of $w_k(\mathbf{v})$ for $\mathbf{v} \in \mathcal{V}$ made above, we can take any fixed set of non-zero vectors $w_1(\mathbf{v}), \dots, w_d(\mathbf{v})$, spanning the tangent cone of $\mathbf{v}$ in $\mathcal{P}$. This does not affect the validity of (1.8) and (1.9).

Theorem 2 implies

Corollary 3. Let $\Delta = \text{conv}(\mathcal{V}) \subset \mathbb{R}^d$ be an arbitrary d -simplex. Then

$$F_{\Delta}(\mathbf{u}) = \frac{d! \text{Vol}(\Delta)}{\prod_{\mathbf{v} \in \mathcal{V}} (1 - \langle \mathbf{v}, \mathbf{u} \rangle)}. \quad (1.10)$$

Remark 2. As we discovered after we proved the above results, statements similar to Corollary 3 in the complex setting can be found in [4, Section 3.5] and in particular [4, Corollary 3.5.6].

A variation of (1.10) also appears in [5], in the context of designing an efficient procedure for integration of polynomials over simplices.

Notice that an arbitrary convex polytope $\mathcal{P}$ admits a triangulation which only uses the existing vertices of $\mathcal{P}$, see e.g. [8, Theorem 3.1]. Applying Corollary 3 and Theorem 1 to the sum of measures corresponding to such a triangulation we get the following.

Corollary 4. The normalized moment generating function $F_{\mathcal{P}}^{\rho}(\mathbf{u})$ of any convex polytope $\mathcal{P}$ with respect to any homogeneous polynomial density function ρ of degree δ is a rational function with denominator dividing

$$\prod_{\mathbf{v} \in \mathcal{V}} (1 - \langle \mathbf{v}, \mathbf{u} \rangle)^{\delta}.$$

Example 1. Let Δ be a triangle in $\mathbb{R}^2$ with vertices $v_1 = (1, 1)$, $v_2 = (2, 5)$ and $v_3 = (3, 2)$. Its normalized moment generating function equals

$$F_{\Delta}(u_1, u_2) = \frac{7}{(1 - u_1 - u_2)(1 - 2u_1 - 5u_2)(1 - 3u_1 - 2u_2)}.$$

Its Taylor expansion about the origin up to the terms of degree 7 is given by

$$\begin{aligned} & 7 + 42u_1 + 56u_2 + 175u_1^2 + 455u_1u_2 + 329u_2^2 + 630u_1^3 + 2387u_1^2u_2 + 3367u_1u_2^2 + 1750u_2^3 + 2107u_1^4 + 10318u_1^3u_2 \\ & + 21217u_1^2u_2^2 + 21546u_1u_2^3 + 8967u_2^4 + 6762u_1^5 + 40082u_1^4u_2 + 106526u_1^3u_2^2 + 157976u_1^2u_2^3 + 128772u_1u_2^4 \\ & + 45276u_2^5 + 21175u_1^6 + 145845u_1^5u_2 + 468895u_1^4u_2^2 + 900123u_1^3u_2^3 + 1074445u_1^2u_2^4 + 741993u_1u_2^5 + 227269u_2^6, \end{aligned}$$

which implies that

$$\begin{aligned} m_{00} &= \frac{7}{2}, m_{10} = 7, m_{01} = \frac{28}{3}, m_{20} = \frac{175}{12}, m_{11} = \frac{455}{24}, m_{02} = \frac{329}{12}, m_{30} = \frac{63}{2}, m_{21} = \frac{2387}{60}, \\ m_{12} &= \frac{3591}{20}, m_{03} = \frac{175}{2}, m_{40} = \frac{2107}{30}, m_{31} = \frac{5159}{60}, m_{22} = \frac{21217}{180}, m_{13} = \frac{3591}{20}, m_{04} = \frac{2989}{10}, \\ m_{50} &= 161, m_{41} = \frac{2863}{15}, m_{32} = \frac{7609}{30}, m_{23} = \frac{5642}{15}, m_{14} = \frac{3066}{5}, m_{05} = 1078, m_{60} = \frac{3025}{8}, \\ m_{51} &= \frac{6945}{16}, m_{42} = \frac{13397}{24}, m_{33} = \frac{128589}{160}, m_{24} = \frac{153493}{120}, m_{15} = \frac{35333}{16}, m_{06} = \frac{32467}{8}. \end{aligned}$$

Results on non-convex polytopes. Our second group of results addresses the problem of distinguishing different polytopes with the same underlying set of vertices from information on their moments. The problem of restoring the vertices of a polygon or a polytope with a constant mass density from information on its moments was addressed earlier in e.g. [12, 15–17, 20, 27]. However, the latter do not provide the recovery of the vertices in the generality required in the present paper. Below we concentrate on the case of constant density and known vertices, and plan to return to the general inverse problem for polytopes with unknown polynomial density and unknown location of their vertices in the future.

First we need to define what we mean by a polytope. It turned out that there is no general consensus about this notion. Instead there exist several competing definitions having their own advantages in different situations. We shall study the following class of polytopal objects.

Definition 1. A subset $\mathcal{P} \subset \mathbb{R}^d$ coinciding with a finite union of arbitrary convex d -dimensional polytopes is called a *generalized polytope*.

Definition 2. The *number of components* of a generalized polytope $\mathcal{P}$ is the number of connected components of the set $\mathcal{P}^\circ \subset \mathcal{P}$ of interior points of $\mathcal{P}$. The closure of each connected component of $\mathcal{P}^\circ$ is called a *component* of $\mathcal{P}$. A generalized polytope with one component is called *indecomposable*.

Remark 3. We say that a simplicial complex in $\mathbb{R}^d$ is *pure* if all its maximal simplices have dimension d . Clearly any generalized polytope in $\mathbb{R}^d$ can be represented as the topological space of an appropriate pure simplicial complex.

Remark 4. Often one considers a more restricted class of objects, namely *polytopes*. A polytope $\mathcal{P} \subset \mathbb{R}^d$ is a generalized polytope homeomorphic to a d -dimensional manifold with boundary.

We need to introduce the notion of a vertex of a generalized polytope.

Definition 3. Given a generalized polytope $\mathcal{P} \subset \mathbb{R}^d$ we call a finite collection of open disjoint d -dimensional simplices in $\mathbb{R}^d$ a *dissection* of $\mathcal{P}$ if the closure of their union coincides with $\mathcal{P}$.

A wealth of material on dissections of polytopes can be found in [21], see also [13].

Definition 4. Given a generalized polytope $\mathcal{P} \subset \mathbb{R}^d$ we call a point $\mathbf{v}$ a *vertex* of $\mathcal{P}$, if $\mathbf{v}$ is a vertex of (the closure of) some open simplex in every dissection of $\mathcal{P}$.

Definition 5. Given a point $p \in \mathcal{P}$ of a generalized polytope $\mathcal{P}$ we denote by the tangent cone $T_p(\mathcal{P})$ of $\mathcal{P}$ at p the set obtained as follows. For a sufficiently small $\epsilon > 0$ set $\mathcal{P}_p(\epsilon) = \mathcal{P} \cap B_p(\epsilon)$ where $B_p(\epsilon)$ is the ϵ -ball centered at p . Define $T_p(\mathcal{P})$ as the set obtained by taking a ray through p and every point of $\mathcal{P}_p(\epsilon)$. In other words, $T_p(\mathcal{P})$ is the cone with the apex at p and the base $B_p(\epsilon)$. (Obviously, $T_p(\mathcal{P})$ is independent of ϵ for a sufficiently small $\epsilon > 0$.)

Lemma 5. A point $\mathbf{v}$ is a *vertex* of $\mathcal{P}$ if and only if $T_{\mathbf{v}}(\mathcal{P})$ does not admit a decomposition in the disjoint union of convex polygonal subcones, such that each subcone in the decomposition has a translation-invariant direction. In particular, if the tangent cone to $\mathcal{P}$ at $\mathbf{v}$ has a connected component with no translation-invariant direction then $\mathbf{v}$ is a vertex.

We denote by $\text{conv}(S)$ the convex hull of an arbitrary set $S \subset \mathbb{R}^d$. The above lemma implies that any vertex of $\text{conv}(\mathcal{P})$ is a vertex of $\mathcal{P}$.

The following result extends Corollary 4 to the case of generalized polytopes.

Proposition 6. For any generalized polytope $\mathcal{P}$ with the set of vertices $\mathcal{V}(\mathcal{P})$, the denominator of its normalized moment generating function $F_{\mathcal{P}}^{\rho}(\mathbf{u})$ with respect to a homogeneous polynomial density function ρ of degree δ divides

$$\Phi_{\mathcal{P}}(\mathbf{u}) := \prod_{\mathbf{v} \in \mathcal{V}(\mathcal{P})} (1 - \langle \mathbf{v}, \mathbf{u} \rangle)^{\delta}.$$

Remark 5. There exist generalized polytopes which do not admit dissections with only existing vertices. The simplest example of this kind is the *Schönhardt polyhedron*, see Figure 1 and [26]. Absence of a dissection $\mathcal{T}$ which uses only its 6 vertices can be established by observing that none of the edges AC , $A'B$, and $B'C'$ can appear in a simplex of $\mathcal{T}$, yet any simplex on these 6 vertices must contain one of them. Therefore, Proposition 6 is not an immediate consequence of Corollary 3.




FIGURE 1. Schönhardt polyhedron obtained from an octahedron (on the left) by removing tetrahedra $[ABB'C']$, $[AA'B'C']$, and $[A'BCC']$.

Remark 6. For “generic” generalized polytopes $\mathcal{P}$ the denominator $\Omega(\mathbf{u})$ of $F_{\mathcal{P}}(\mathbf{u})$ equals $\Phi_{\mathcal{P}}(\mathbf{u})$, but for certain special polytopes the denominator $\Omega(\mathbf{u})$ may be its proper divisor, as can be seen from the following example. Let $A = \{0, a_1, a_2, a_3\} \subset \mathbb{R}^3$ be a spanning set, and $v \in \mathbb{R}^3$. Let $\mathcal{P}_{\pm} := \text{conv}(v \pm A)$ and $\mathcal{P} := \mathcal{P}_+ \cup \mathcal{P}_-$. Then $1 - \langle \mathbf{u}, v \rangle$ does not appear in $\Omega(\mathbf{u})$, as

$$F_{\mathcal{P}}(\mathbf{u}) = F_{\mathcal{P}_+}(\mathbf{u}) + F_{\mathcal{P}_-}(\mathbf{u}) = K \frac{\sum_{1 \leq i < j \leq 3} \langle \mathbf{u}, a_i \rangle \langle \mathbf{u}, a_j \rangle + (1 - \langle \mathbf{u}, v \rangle)^2}{\prod_{1 \leq i \leq 3} ((1 - \langle \mathbf{u}, v \rangle)^2 - \langle \mathbf{u}, a_i \rangle^2)},$$

where $K \neq 0$ is a real constant.

Now we introduce several finite-dimensional linear spaces related to a given finite spanning set $S \subset \mathbb{R}^d$. Let $\mathcal{P}(S)$ be the set of all generalized polytopes $\mathcal{P}$ whose sets $\mathcal{V}(\mathcal{P})$ of vertices are contained in S . For $\mathcal{P} \in \mathcal{P}(S)$ we denote by $\mu_{\mathcal{P}}$ its standard measure. (Obviously, $\mu_{\mathcal{P}}$ is supported on $\mathcal{P} \subseteq \text{conv}(S)$.)

Denote by $\mathfrak{M}(S)$ the linear space of all signed measures, i.e. the linear span of all standard measures $\mu_{\mathcal{P}}$ for $\mathcal{P} \in \mathcal{P}(S)$. Let $\mathfrak{M}^{\Delta}(S) \subseteq \mathfrak{M}(S)$ be its subspace spanned by μ_{Δ} , for $\Delta \in \mathcal{P}(S)$ a d -dimensional simplex. (The space $\mathfrak{M}^{\Delta}(S)$ has earlier appeared in [1], [2], [3] in a somewhat different context.) We shall refer to elements of $\mathfrak{M}(S)$ as to *polytopal measures* with the vertex set S . The next conjecture is central to our study.

Conjecture 7. For an arbitrary spanning set S and any generalized polytope $\mathcal{P}$ with set of vertices contained in S , its standard measure $\mu_{\mathcal{P}}$ belongs to $\mathfrak{M}^{\Delta}(S)$. In other words, $\mathfrak{M}(S) = \mathfrak{M}^{\Delta}(S)$.

By Remark 5, the above conjecture is non-trivial. While we do not have a proof of Conjecture 7 in its full generality, we have succeeded in proving it for a rather large class of spanning sets. Roughly speaking, the latter should be close to “generic”. Specifically, given a finite spanning set $S \subset \mathbb{R}^d$, we say that S is *weakly non-degenerate* if any $(d+2)$ -tuple of points from S is spanning. If S satisfies the stronger condition that each $(d+1)$ -subset of S is spanning then we call the latter S *strongly non-degenerate*.

Theorem 8. Conjecture 7 holds for any weakly non-degenerate finite set S .

Remark 7. Theorem 8 would imply Conjecture 7 if one could prove that the standard measure of an arbitrary generalized polytope $\mathcal{P}$ can be obtained as the limit of the standard measures of a 1-parameter family of generalized polytopes $\mathcal{P}(t)$ with $\mathcal{P}(0) = \mathcal{P}$ such that for $t \neq 0$ the vertices of $\mathcal{P}(t)$ are weakly non-degenerate, and each vertex of $\mathcal{P}(t)$ tending to a vertex of $\mathcal{P}$ as $t \rightarrow 0$. Unfortunately we are unable to prove the existence of such deformations in general.

The key idea in the proof of Theorem 8 is to study the corresponding spaces of Fantappiè transformations of signed measures in $\mathfrak{M}(S)$. In particular, we are able to compute the corresponding dimensions¹. In more detail, let $\mathfrak{F}(S)$ (resp. $\mathfrak{F}^{\Delta}(S)$) be the linear space of Fantappiè transformations of signed measures in $\mathfrak{M}(S)$ (resp. $\mathfrak{M}^{\Delta}(S)$). In other words, $\mathfrak{F}(S)$ (resp. $\mathfrak{F}^{\Delta}(S)$) is the space of normalized moment generating functions of signed measures in $\mathfrak{M}(S)$ (resp. $\mathfrak{M}^{\Delta}(S)$).

Since each compactly supported measure is uniquely determined by its complete set of moments, the map

$$F_{\mu} : \mathfrak{M}(S) \rightarrow \mathfrak{F}(S), \quad (1.11)$$

induced by the Fantappiè transformation is a linear isomorphism, cf. [4, Section 3.5].

Finally, given a spanning set $S = \{\mathbf{v}_1, \dots, \mathbf{v}_N\} \subset \mathbb{R}^d$, denote by $\mathfrak{Rat}(S)$ the linear space of all rational functions with the denominator $\Phi_S(\mathbf{u})$ as in (1.12),

$$\Phi_S(\mathbf{u}) = \prod_{i=1}^N (1 - \langle \mathbf{v}_i, \mathbf{u} \rangle), \quad (1.12)$$

and with the numerator an arbitrary real (inhomogeneous) polynomial of degree at most $N - d - 1$. Here the numerator and the denominator might have common factors.

Proposition 9. $\mathfrak{F}^{\Delta}(S)$ coincides with $\mathfrak{Rat}(S)$ if and only if S is strongly non-degenerate.

Corollary 10. If S is strongly non-degenerate then $\mathfrak{M}^{\Delta}(S) = \mathfrak{M}(S)$.

Corollary 10 implies that for strongly non-degenerate S the dimension of all these linear spaces equals $\binom{N-1}{d}$. Note that Corollary 10 settles Theorem 8 for the strongly non-degenerate S .

Our final goal is to explicitly solve the following inverse moment problem.

¹ Note that presently we are not aware of a formula or a recipe for calculating the dimension of $\mathfrak{M}^{\Delta}(S)$ without the assumption of the theorem. (The manuscript [2] contains an algorithm constructing a basis of this space.)

Problem 1. Given a strongly non-degenerate spanning set $S \subset \mathbb{R}^d$, $|S| = N$, find the unique polytopal measure in $\mathfrak{M}(S)$ with a given set of all moments up to order $N - d - 1$.

We start with the following simple observation.

Lemma 11. Given an arbitrary spanning set $S \subset \mathbb{R}^d$, $|S| = N$, and an arbitrary polynomial $T(\mathbf{u})$ of degree at most $N - d - 1$ there exists a unique rational function $R(\mathbf{u}) = P(\mathbf{u})/\Phi_S(\mathbf{u})$ with Taylor polynomial of degree $N - d - 1$ at the origin equal to $T(\mathbf{u})$. Namely, $P(\mathbf{u}) = [T(\mathbf{u})\Phi_S(\mathbf{u})]_{N-d-1}$, where $[\cdot]_{N-d-1}$ stands for the truncated polynomial with all monomials up to degree $N - d - 1$.

For $S = \{\mathbf{v}_1, \dots, \mathbf{v}_N\} \subset \mathbb{R}^d$ strongly non-degenerate, we give an explicit inversion formula determining the densities of an unknown polytopal measure having a given set of moments up to order $N - d - 1$ on each simplex in a natural basis of $\mathfrak{M}^\Delta(S)$. In view of Lemma 11 we can assume that we are already given an arbitrary rational function $R(\mathbf{u}) = P(\mathbf{u})/\Phi_S(\mathbf{u})$, where $\deg P(\mathbf{u}) \leq N - d - 1$, and we want to determine the densities of the required signed measure from $\mathfrak{M}(S)$ in terms of numerator $P(\mathbf{u})$.

From now on we shall choose the basis of $\mathfrak{M}^\Delta(S)$ consisting of the standard measures of all simplices containing the last vertex $\mathbf{v}_N$, see Lemma 15 below. Let $\mathfrak{L} = \{l_1, l_2, \dots, l_{N-1}\}$ be the $(N - 1)$ -tuple of linear forms corresponding to vertices $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{N-1}$, where $l_i(\mathbf{u}) = 1 - \langle \mathbf{v}_i, \mathbf{u} \rangle$. Consider the linear span $V_{\mathfrak{L}}$ of all possible products of the form $l_{j_1} \cdot l_{j_2} \cdot \dots \cdot l_{j_{N-d-1}}$, $1 \leq j_1 < j_2 < \dots < j_{N-d-1}$. There are $\binom{N-1}{d}$ such products, and each of them is a polynomial of degree at most $N - d - 1$. On the other hand, the dimension of the space $Pol(N - d - 1, d)$ of all (inhomogeneous) polynomials of degree at most $N - d - 1$ in d variables equals $\binom{N-1}{d}$, as well.

Define the square matrix Mat_S of size $\binom{N-1}{d}$ with entries being coefficients of the above products of linear forms with respect to the standard monomial basis in $Pol(N - d - 1, d)$. We assume that Mat_S acts on the space $V_{\mathfrak{L}}$ of column vectors.

Theorem 12. For an arbitrary strongly non-degenerate spanning set $S \subset \mathbb{R}^d$, $|S| = N$, the matrix Mat_S is invertible. Moreover, for a rational function $R(\mathbf{u}) = P(\mathbf{u})/\Phi_S(\mathbf{u})$, where $P(\mathbf{u})$ is an arbitrary polynomial of degree $N - d - 1$, there exists a unique measure $\mu_R \in \mathfrak{M}(S)$ with Fantappiè transform $R(\mathbf{u})$. Namely,

$$\mu_R = Mat_S^{-1}(P(\mathbf{u})). \quad (1.13)$$

Remark 8. A detailed explanation of the meaning of (1.13) can be found in the proof of Theorem 12, see also Example 2 below. An explicit formula for the matrix Mat_S^{-1} is given in Lemma 18.

Recall that a spanning set S is weakly non-degenerate if any $(d + 2)$ -tuple of its points is spanning. With minor changes, the above solution of the inverse moment problem can be adapted to this more general case. In order not to overload the introduction we refer the readers interested in this situation to Section 4. The case of an arbitrary spanning set S , however, remains unsolved and offers several interesting challenges in matroid theory. We hope to return to it in the future.

It will be convenient to work with scaled volumes of simplices, which we call *weights*.

Definition 6. Given a signed measure μ in $\mathbb{R}^d$ and a d -dimensional simplex $\Delta \subset \mathbb{R}^d$ we define the *weight* w_Δ of Δ by the formula:

$$w_\Delta = d! \int_{\Delta} d\mu. \quad (1.14)$$

In other words, the density d_Δ of the measure in question which should be placed at Δ equals

$$d_\Delta = \frac{w_\Delta}{d! \text{Vol}(\Delta)}.$$

We finish the introduction by explicitly solving the above inverse problem for a concrete 5-tuple of points in $\mathbb{R}^2$.

Example 2. Set $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4, \mathbf{v}_5\}$ where $\mathbf{v}_1 = (1, 0)$, $\mathbf{v}_2 = (2, 1)$, $\mathbf{v}_3 = (1, 2)$, $\mathbf{v}_4 = (0, 1)$, $\mathbf{v}_5 = (0, 0)$. The corresponding set $\mathfrak{L} = \{l_1, l_2, l_3, l_4\}$ of linear forms is given by $l_1 = 1 - u_1$, $l_2 = 1 - 2u_1 - u_2$, $l_3 = 1 - u_1 - 2u_2$, $l_4 = 1 - u_2$. Additionally, $l_5 = 1$. We are considering the basis of $\mathfrak{M}^\Delta(S)$ consisting of (the standard measures of) 6 triangles containing $\mathbf{v}_5$. Therefore we need 6 quadratic forms obtained as pairwise products $l_i l_j$, $1 \leq i < j \leq 4$. We get

$$\begin{cases} l_1 l_2 = 1 - 3u_1 - u_2 + 2u_1^2 + u_1 u_2 \\ l_1 l_3 = 1 - 2u_1 - 2u_2 + u_1^2 + 2u_1 u_2 \\ l_1 l_4 = 1 - u_1 - u_2 + u_1 u_2 \\ l_2 l_3 = 1 - 3u_1 - 3u_2 + 2u_1^2 + 5u_1 u_2 + 2u_2^2 \\ l_2 l_4 = 1 - 2u_1 - 2u_2 + 2u_1 u_2 + u_2^2 \\ l_3 l_4 = 1 - u_1 - 3u_2 + u_1 u_2 + 2u_2^2. \end{cases}$$

Notice that $l_1 l_2$ corresponds to triangle Δ_{345} , $l_1 l_3$ to Δ_{245} , $l_1 l_4$ to Δ_{234} , $l_2 l_3$ to Δ_{145} , $l_2 l_4$ to Δ_{135} , and $l_3 l_4$ to Δ_{125} . Ordering monomials spanning the space $\text{Pol}(2, 2)$ as $(1, u_1, u_2, u_1^2, u_1 u_2, u_2^2)$ we get the 6×6 -matrix Mat_S and its inverse Mat_S^{-1} as follows

$$\begin{array}{c} \text{Mat}_S = \\ \begin{matrix} & l_1 l_2 & l_1 l_3 & l_1 l_4 & l_2 l_3 & l_2 l_4 & l_3 l_4 \end{matrix} \\ \begin{matrix} 1 \\ u_1 \\ u_2 \\ u_1^2 \\ u_1 u_2 \\ u_2^2 \end{matrix} \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ -3 & -2 & -1 & -3 & -2 & -1 \\ -1 & -2 & -1 & -3 & -2 & -3 \\ 2 & 1 & 0 & 2 & 0 & 0 \\ 1 & 2 & 1 & 5 & 2 & 1 \\ 0 & 0 & 0 & 2 & 1 & 2 \end{pmatrix} \end{array} \quad \begin{array}{c} 4\text{Mat}_S^{-1} = \\ \begin{matrix} 1 & u_1 & u_2 & u_1^2 & u_1 u_2 & u_2^2 \end{matrix} \\ \begin{pmatrix} 1 & -1 & 1 & 1 & 1 & -1 \\ -4 & 0 & -4 & 0 & 0 & -4 \\ 9 & 3 & 3 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \\ -4 & -4 & 0 & -4 & 0 & 0 \\ 1 & 1 & -1 & 1 & -1 & 1 \end{pmatrix} \end{array}$$

(For $\text{T}_\text{E}\text{X}$ reasons we give 4Mat_S^{-1} above.) Thus, given an arbitrary rational function $R(u_1, u_2) = P(u_1, u_2)/\Phi_S(u_1, u_2)$ where $P(u_1, u_2) = a_{00} + a_{1,0}u_1 + a_{0,1}u_2 + a_{2,0}u_1^2 + a_{1,1}u_1 u_2 + a_{0,2}u_2^2$ is a polynomial of degree at most 2 and $\Phi_S(u_1, u_2) = l_1 l_2 l_3 l_4 l_5$, we get

$$\begin{cases} w_{345} = \frac{1}{4}(a_{00} - a_{10} + a_{01} + a_{20} - a_{11} + a_{02}) \\ w_{245} = -a_{00} - a_{01} - a_{02} \\ w_{235} = \frac{1}{4}(9a_{00} + 3a_{01} + 3a_{10} + a_{20} + a_{11} + a_{02}) \\ w_{145} = \frac{1}{4}(a_{00} + a_{01} + a_{10} + a_{20} + a_{11} + a_{02}) \\ w_{135} = -a_{00} - a_{10} - a_{20} \\ w_{125} = \frac{1}{4}(a_{00} + a_{10} - a_{01} + a_{20} - a_{11} + a_{02}), \end{cases}$$

where w_{ijk} is the weight of the signed measure to be placed on Δ_{ijk} , see (1.14).

To illustrate all steps of solution of our inverse moment problem assume that we are looking for a polygonal measure with the vertex set S and (ad hoc chosen) moments $m_{00} = 1, m_{10} = 2, m_{01} = 3, m_{20} = 4, m_{11} = 5, m_{02} = 6$. Then its

normalized moment generating function $F_\mu(\mathbf{u})$ satisfies the relation

$$F_\mu(\mathbf{u}) = 1 + \frac{2!}{0!0!} + 2\frac{3!}{1!0!}u_1 + 3\frac{3!}{0!1!}u_2 + 4\frac{4!}{2!0!}u_1^2 + 5\frac{4!}{1!1!}u_1u_2 + 6\frac{4!}{0!2!}u_2^2 + \dots = \frac{P(u_1, u_2)}{l_1l_2l_3l_4l_5},$$

where $P(u_1, u_2)$ is a (non-homogeneous) polynomial of at most second degree. Thus, truncating the product of the left-hand side and $l_1l_2l_3l_4l_5$ up to the second degree, we obtain

$$P(u_1, u_2) = 2 + 4u_1 + 10u_2 + 10u_1^2 + 24u_1u_2 + 10u_2^2,$$

i.e. $a_{00} = 2, a_{10} = 4, a_{01} = 10, a_{20} = 10, a_{11} = 24, a_{02} = 10$. Thus $w_{345} = 1, w_{245} = -22, w_{235} = 26, w_{145} = 15, w_{135} = -16, w_{125} = -2$. The areas of the corresponding triangles are equal to: $\text{Area}(\Delta_{345}) = \frac{1}{2}; \text{Area}(\Delta_{245}) = 1; \text{Area}(\Delta_{235}) = \frac{3}{2}; \text{Area}(\Delta_{145}) = \frac{1}{2}; \text{Area}(\Delta_{135}) = 1; \text{Area}(\Delta_{125}) = \frac{1}{2}$. This implies that the densities of the measure of the corresponding triangles are equal to $d_{345} = 1, d_{245} = -11, d_{235} = \frac{26}{3}, d_{145} = 15, d_{135} = -8, d_{125} = -2$. To obtain the final densities in the convex hull $\text{conv}(S)$ of S one has to decompose $\text{conv}(S)$ into domains obtained by removing from $\text{conv}(S)$ the set of all hyperplanes spanned by vertices in S . For each such domain we should add up the densities of all basic simplices containing this domain. The resulting measure is shown in Fig. 2.




FIGURE 2. Final measure in Example 2.

Remark 9. Domains into which the convex hull $\text{conv}(S)$ is cut by the hyperplanes spanned by S were introduced in [1] where they were called *chambers*. The incidence matrix of the simplices spanned by S and those chambers was studied in some detail in [2],[3]. This matrix allows to formalize the last step of construction of the above polygonal measure, where information on the densities of the simplices is transformed into information on the densities of the chambers. But, in general, already the number of chambers is a complicated invariant of the set S . It seems that the general problem of constructing the set of chambers and the corresponding incidence matrix in terms of a given S is quite non-trivial.

Acknowledgement. The second author is grateful to the Mathematics Department of Stockholm University for the hospitality in June 2011 when this project was initiated. The third author wants to acknowledge the hospitality of the School of Physical and Mathematical Sciences, Nanyang Technological University in April 2012 when this project was completed. We want to thank Sinai Robins for numerous discussions of the topic. We acknowledge extremely helpful answers and comments on our questions on mathoverflow.net, in particular ones by David Eppstein, Dirk Lorenz, Igor Pak, David Speyer, and Gjergji Zaimi. Finally, the third author wants to thank late Mikael Passare (who unfortunately left us so early) for discussions of the properties of Fantappiè transformation and for pointing out reference [4] in September 2011.

2. PROVING RESULTS ON CONVEX POLYTOPES

Following Brion-Lawrence-Khovanskii-Pukhlikov-Barvinok, see [6, 8, 16, 18], we define for each vector $\mathbf{z} \in \mathbb{R}^d$ the j -th *axial* moment $\mu_j(\mathbf{z})$ of a simple convex polytope $\mathcal{P}$ with respect to $\mathbf{z}$ as

$$\mu_j(\mathbf{z}) = \int_{\mathcal{P}} \langle \mathbf{x}, \mathbf{z} \rangle^j d\mathbf{x}.$$

We will use the following important statement, cf. e.g. [8, Theorem 10.5].

Theorem 13. The moment $\mu_j(\mathbf{z})$ satisfies

$$\mu_j(\mathbf{z}) = \frac{(-1)^d j!}{(j+d)!} \sum_{\mathbf{v} \in \mathcal{V}} \langle \mathbf{v}, \mathbf{z} \rangle^{j+d} D_{\mathbf{v}}(\mathbf{z}), \quad (2.1)$$

where $D_{\mathbf{v}}(\mathbf{z}) := \frac{|\det K_{\mathbf{v}}|}{\prod_{j=1}^d \langle w_j(\mathbf{v}), \mathbf{z} \rangle}$, and $\mathbf{z}$ is an arbitrary vector for which the products $\prod_{j=1}^d \langle w_j(\mathbf{v}), \mathbf{z} \rangle$, $\mathbf{v} \in \mathcal{V}$, do not vanish. Moreover, the following identities hold:

$$\sum_{\mathbf{v} \in \mathcal{V}} \langle \mathbf{v}, \mathbf{z} \rangle^j D_{\mathbf{v}}(\mathbf{z}) = 0, \quad \text{for } 0 \leq j \leq d-1. \quad (2.2)$$

Proof of Theorem 2. To prove (1.8), consider the generating function

$$\Psi_{\mathbf{z}}(u) = \sum_{j=0}^{\infty} \frac{(j+d)!}{j!} \mu_j(\mathbf{z}) u^j,$$

where $u \in \mathbb{R}$. Formula (2.1) implies that $\Psi_{\mathbf{z}}(u)$ is rational. Indeed,

$$\begin{aligned} \Psi_{\mathbf{z}}(u) &= \sum_{j=0}^{\infty} (-1)^d \sum_{\mathbf{v} \in \mathcal{V}} \langle \mathbf{v}, \mathbf{z} \rangle^{j+d} \frac{|\det K_{\mathbf{v}}| u^j}{\prod_{k=1}^d \langle w_k(\mathbf{v}), \mathbf{z} \rangle} = \\ &= (-1)^d \sum_{\mathbf{v} \in \mathcal{V}} \frac{\langle \mathbf{v}, \mathbf{z} \rangle^d |\det K_{\mathbf{v}}|}{\prod_{k=1}^d \langle w_k(\mathbf{v}), \mathbf{z} \rangle} \sum_{j=0}^{\infty} \langle \mathbf{v}, \mathbf{z} \rangle^j u^j = \sum_{\mathbf{v} \in \mathcal{V}} \frac{\langle \mathbf{v}, \mathbf{u} \rangle^d |\det K_{\mathbf{v}}|}{\prod_{k=1}^d \langle w_k(\mathbf{v}), \mathbf{u} \rangle} \cdot \frac{(-1)^d}{1 - \langle \mathbf{v}, \mathbf{u} \rangle}, \end{aligned}$$

where $\mathbf{u} = u\mathbf{z}$. On the other hand, using the multinomial coefficients $\binom{|J|}{J} = \frac{|J|!}{j_1! \dots j_d!}$ of multiindices $J = (j_1, \dots, j_d) \vdash |J|$, one gets

$$\int_{\mathcal{P}} \langle \mathbf{x}, \mathbf{z} \rangle^j d\mathbf{x} = \int_{\mathcal{P}} \left(\sum_{i=1}^d x_i z_i \right)^j d\mathbf{x} = \sum_{J \vdash j} \binom{j}{J} \mathbf{z}^J \int_{\mathcal{P}} \mathbf{x}^J d\mathbf{x} = \sum_{J \vdash j} \binom{j}{J} \mathbf{z}^J m_j(\mathcal{P}),$$

where $m_J(\mathcal{P}) = m_J(\mu_{\mathcal{P}})$. Therefore,

$$\begin{aligned} F_{\mathcal{P}}(u\mathbf{z}) &:= F_{\mathcal{P}}(uz_1, \dots, uz_d) := \sum_{j=0}^{\infty} \sum_{(j_1, \dots, j_d) \vdash j} \frac{(j+d)!}{j_1! \dots j_d!} m_{j_1, \dots, j_d}(uz_1)^{j_1} \dots (uz_d)^{j_d} \\ &= \sum_{j=0}^{\infty} \frac{(j+d)!}{j!} \left[\sum_{J:=(j_1, \dots, j_d) \vdash j} \binom{j}{J} m_J(\mathcal{P}) \mathbf{z}^J \right] u^j = \sum_{j=0}^{\infty} \frac{(j+d)!}{j!} \mu_j(\mathbf{z}) u^j = \Psi_{\mathbf{z}}(u), \end{aligned}$$

and (1.8) follows.

In view of relations (2.2) the right-hand side of (1.8) can be rewritten as (1.9). Indeed, writing $(1 - \langle \mathbf{v}, \mathbf{u} \rangle)^{-1} = \sum_{j=0}^{\infty} \langle \mathbf{v}, \mathbf{u} \rangle^j$ and expanding (1.8) with respect to j th powers of $\langle \mathbf{v}, \mathbf{u} \rangle$, we see that (2.2) implies that for $j < d$ the sum of all terms $\langle \mathbf{v}, \mathbf{u} \rangle^j$ vanishes. $\square$

Proof of Corollary 3. Let $\mathcal{V} = (\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_d)$. Then for each $j \neq i$ we have $w_j(\mathbf{v}_i) = \mathbf{v}_j - \mathbf{v}_i$. Hence $|\det K_{\mathbf{v}_i}|$ does not depend upon i and equals $d! \text{Vol}(\Delta)$. The right-hand side of (1.8) becomes

$$\begin{aligned} (-1)^d d! \text{Vol}(\Delta) \sum_{i=0}^d \frac{\langle \mathbf{v}_i, \mathbf{u} \rangle^d}{\prod_{j=1}^d \langle w_j(\mathbf{v}_i), \mathbf{u} \rangle} \cdot \frac{1}{1 - \langle \mathbf{v}_i, \mathbf{u} \rangle} &= \\ &= (-1)^d d! \text{Vol}(\Delta) \sum_{i=0}^d \frac{\langle \mathbf{v}_i, \mathbf{u} \rangle^d}{\prod_{j \neq i} \langle \mathbf{v}_j - \mathbf{v}_i, \mathbf{u} \rangle} \cdot \frac{1}{1 - \langle \mathbf{v}_i, \mathbf{u} \rangle} = \\ &= (-1)^d d! \text{Vol}(\Delta) \sum_{i=0}^d \frac{\zeta_i^d}{\prod_{j \neq i} (\zeta_j - \zeta_i)} \cdot \frac{1}{1 - \zeta_i}, \end{aligned}$$

where $\zeta_i = \langle \mathbf{v}_i, \mathbf{u} \rangle$. Computing the common denominator of the latter, we obtain

$$F_{\Delta}(\mathbf{u}) = \frac{(-1)^d d! \text{Vol}(\Delta)}{\prod_{i=0}^d (1 - \zeta_i)} \frac{\sum_{i=0}^d \left[\prod_{k>l, k \neq i \neq l} (\zeta_k - \zeta_l) \prod_{j \neq i} (1 - \zeta_j) \right] (-1)^i \zeta_i^d}{\prod_{s>t} (\zeta_s - \zeta_t)}.$$

It is convenient to introduce one more linear form $\zeta_{d+1} := 1$, so that the last expression reads as

$$F_{\Delta}(\mathbf{u}) = (-1)^d d! \text{Vol}(\Delta) \frac{\sum_{i=0}^d \left[\prod_{\substack{d+1 \geq k > l \geq 0, \\ k \neq i \neq l}} (\zeta_k - \zeta_l) \right] (-1)^i \zeta_i^d}{\prod_{d+1 \geq s > t \geq 0} (\zeta_s - \zeta_t)}. \quad (2.3)$$

To complete the proof we notice that

$$\begin{aligned}
0 &= \det \begin{pmatrix} 1 & 1 & \dots & 1 & 1 \\ \zeta_0 & \zeta_1 & \dots & \zeta_d & 1 \\ \zeta_0^2 & \zeta_1^2 & \dots & \zeta_d^2 & 1 \\ \dots & \dots & \dots & \dots & \dots \\ \zeta_0^d & \zeta_1^d & \dots & \zeta_d^d & 1 \\ \zeta_0^d & \zeta_1^d & \dots & \zeta_d^d & 1 \end{pmatrix} = 1 \cdot \det \begin{pmatrix} 1 & 1 & \dots & 1 \\ \zeta_0 & \zeta_1 & \dots & \zeta_d \\ \zeta_0^2 & \zeta_1^2 & \dots & \zeta_d^2 \\ \dots & \dots & \dots & \dots \\ \zeta_0^d & \zeta_1^d & \dots & \zeta_d^d \end{pmatrix} + \\
&+ (-1)^{d+1} \sum_{i=0}^d \zeta_i^d (-1)^i \cdot \det \begin{pmatrix} 1 & \dots & 1 & 1 & \dots & 1 \\ \zeta_0 & \dots & \zeta_{i-1} & \zeta_{i+1} & \dots & \zeta_{d+1} \\ \zeta_0^2 & \dots & \zeta_{i-1}^2 & \zeta_{i+1}^2 & \dots & \zeta_{d+1}^2 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \zeta_0^d & \dots & \zeta_{i-1}^d & \zeta_{i+1}^d & \dots & \zeta_{d+1}^d \end{pmatrix} = \\
&= \prod_{d \geq k > l \geq 0} (\zeta_k - \zeta_l) + (-1)^{d+1} \sum_{i=0}^d \zeta_i^d (-1)^i \cdot \left[\prod_{\substack{d+1 \geq k > l \geq 0, \\ k \neq i \neq l}} (\zeta_k - \zeta_l) \right].
\end{aligned}$$

Indeed, the first matrix has two identical rows and thus vanishing determinant, which we expand with respect to the last row. The last equality is the standard formula for the Vandermonde determinant. Thus we have

$$\sum_{i=0}^d \zeta_i^d (-1)^i \cdot \left[\prod_{\substack{d+1 \geq k > l \geq 0, \\ k \neq i \neq l}} (\zeta_k - \zeta_l) \right] = (-1)^d \prod_{d \geq k > l \geq 0} (\zeta_k - \zeta_l).$$

Now we plug this formula into (2.3) and get

$$F_\Delta(\mathbf{u}) = d! \text{Vol}(\Delta) \frac{1}{\prod_{t=0}^d (1 - \zeta_t)} = \frac{d! \text{Vol}(\Delta)}{\prod_{\mathbf{v} \in \mathcal{V}} (1 - \langle \mathbf{v}, \mathbf{u} \rangle)}. \quad \square$$

Lemma 14. Let $\mathbf{u} = (u_1, \dots, u_d)$ and $\mathbf{x} = (x_1, \dots, x_d)$ be formal variables, and $\ell \in \mathbb{R}$. Then

$$\left(\sum_k u_k \frac{\partial}{\partial u_k} + \ell \right) \circ (1 - \langle \mathbf{x}, \mathbf{u} \rangle)^{-\ell} = \ell (1 - \langle \mathbf{x}, \mathbf{u} \rangle)^{-\ell-1}. \quad (2.4)$$

Proof. Note that $u_k \frac{\partial}{\partial u_k} \circ (1 - \langle \mathbf{x}, \mathbf{u} \rangle)^{-\ell} = x_k u_k \ell (1 - \langle \mathbf{x}, \mathbf{u} \rangle)^{-\ell-1}$. Thus

$$\begin{aligned}
\left(\sum_k u_k \frac{\partial}{\partial u_k} + \ell \right) \circ (1 - \langle \mathbf{x}, \mathbf{u} \rangle)^{-\ell} &= \ell (1 - \langle \mathbf{x}, \mathbf{u} \rangle)^{-\ell-1} \langle \mathbf{x}, \mathbf{u} \rangle + \ell (1 - \langle \mathbf{x}, \mathbf{u} \rangle)^{-\ell} \\
&= \ell (1 - \langle \mathbf{x}, \mathbf{u} \rangle)^{-\ell-1}. \quad \square
\end{aligned}$$

Proof of (1.3). For a d -variate polynomial $g(\mathbf{z})$ we denote by $g(\mathbf{u} \frac{\partial}{\partial \mathbf{u}})$ the differential operator $g(u_1 \frac{\partial}{\partial u_1}, \dots, u_d \frac{\partial}{\partial u_d})$. We use the identity

$$g\left(\mathbf{u} \frac{\partial}{\partial \mathbf{u}}\right) \circ \sum_I a_I \mathbf{x}^I \mathbf{u}^I = \sum_{I \geq 0} g(I) a_I \mathbf{x}^I \mathbf{u}^I, \quad (2.5)$$

which holds for any formal d -variate power series $\sum_I a_I \mathbf{x}^I \mathbf{u}^I$ and any d -variate polynomial $g(\mathbf{z})$. (It can be easily verified for monomial $g(\mathbf{z})$ and then extended by

linearity.) Setting $h(\mathbf{z}) := \prod_{\ell=1}^d \left(\sum_{k=1}^d z_k + \ell \right)$, notice that $h(I) = (|I| + 1)(|I| + 2) \cdots (|I| + d)$. Now using (2.5) together with the obvious identity:

$$(1 - \langle \mathbf{x}, \mathbf{u} \rangle)^{-1} = \sum_{I \geq 0} \binom{|I|}{I} \mathbf{x}^I \mathbf{u}^I,$$

one obtains

$$\begin{aligned} F_\mu(\mathbf{u}) &:= \sum_{I \geq 0} \binom{|I| + d}{I} m_I(\mu) \mathbf{u}^I = \sum_{I \geq 0} h(I) \binom{|I|}{I} m_I(\mu) \mathbf{u}^I \\ &= \int_{\mathbb{R}^d} \sum_{I \geq 0} h(I) \binom{|I|}{I} \mathbf{x}^I \mathbf{u}^I d\mu(\mathbf{x}) = \int_{\mathbb{R}^d} h\left(\mathbf{u} \frac{\partial}{\partial \mathbf{u}}\right) \circ \sum_{I \geq 0} \binom{|I|}{I} \mathbf{x}^I \mathbf{u}^I d\mu(\mathbf{x}) \\ &= \int_{\mathbb{R}^d} h\left(\mathbf{u} \frac{\partial}{\partial \mathbf{u}}\right) \circ \frac{d\mu(\mathbf{x})}{1 - \langle \mathbf{x}, \mathbf{u} \rangle} = \int_{\mathbb{R}^d} \frac{d! \, d\mu(\mathbf{x})}{(1 - \langle \mathbf{x}, \mathbf{u} \rangle)^{d+1}}, \end{aligned}$$

where in the final derivation we repeatedly made use of (2.4), for $1 \leq \ell \leq d$. $\square$

Remark 10. Another point of view on (1.3) is that it is the result of the application of the differential operator $g\left(\mathbf{u} \frac{\partial}{\partial \mathbf{u}}\right)$ to the integral transformation $\int_{\mathbb{R}^d} \frac{d\mu(\mathbf{x})}{1 - \langle \mathbf{x}, \mathbf{u} \rangle}$ of the measure μ (also known as the Fantappiè transform of μ); see e.g. [19].

In [22] a similar idea was applied to the harmonic polygonal measures in the plane.

Proof of Theorem 1. Assume first that $\rho(x_1, \dots, x_d) = \mathbf{x}^K = x_1^{k_1} \cdots x_d^{k_d}$ is a monomial and consider

$$\rho\left(\frac{\partial}{\partial \mathbf{u}}\right) \circ F_\mu(\mathbf{u}) = \frac{\partial^{|K|}}{\partial u_1^{k_1} \cdots \partial u_d^{k_d}} \circ \sum_{I=(i_1, \dots, i_d) \geq 0} \frac{(|I| + d)!}{i_1! \cdots i_d!} m_I(\mu) \mathbf{u}^I.$$

One gets

$$\begin{aligned} \frac{\partial^{|K|}}{\partial \mathbf{u}^K} \circ F_\mu(\mathbf{u}) &= \sum_{I=(i_1, \dots, i_d) \geq 0} \left(\frac{(|I| + |K| + d)!}{\prod_{j=1}^d (i_j + k_j)!} \prod_{j=1}^d \frac{(i_j + k_j)!}{i_j!} \right) m_{I+K}(\mu) \mathbf{u}^I \\ &= \sum_{I=(i_1, \dots, i_d) \geq 0} \frac{(|I| + d + |K|)!}{\prod_{j=1}^d i_j!} m_I(\mathbf{x}^K \mu) \mathbf{u}^I. \end{aligned}$$

Observe that the normalizing coefficients of $m_I(\mathbf{x}^K \mu)$ in the latter expression depend only on I and $|K|$ but not on particular entries of K . Therefore for an arbitrary homogeneous ρ of degree δ one gets by additivity

$$\rho\left(\frac{\partial}{\partial \mathbf{u}}\right) \circ F_\mu(\mathbf{u}) = \sum_{I=(i_1, \dots, i_d) \geq 0} \frac{(|I| + d + \delta)!}{\prod_{j=1}^d i_j!} m_I(\rho\mu).$$

This shows (1.6). Repeated application of (2.4), for $d + 1 \leq \ell \leq d + \delta$, to the integral representation (1.3), respectively, to the representation (1.2), of $F_{\rho\mu}(\mathbf{u})$ implies (1.7), respectively, (1.5). $\square$

3. INVERSE MOMENT PROBLEM FOR STRONGLY NON-DEGENERATE S

Proof of Lemma 5. We prove first that the tangent cone at any non-vertex allows a decomposition into convex polytopal cones each having a translation-invariant direction.

Let $\mathbf{v}$ be a point in $\mathcal{P}$ which is not a vertex. Then there is a dissection $\mathcal{T}$ of $\mathcal{P}$ such that $\mathbf{v}$ is not a vertex of any simplex of $\mathcal{T}$. Let U be the set of simplices S_u of $\mathcal{T}$ with closures containing $\mathbf{v}$. Take the dissection of the tangent cone $T_{\mathbf{v}}(\mathcal{P})$ into the tangent cones to simplices from U , $T_{\mathbf{v}}(\mathcal{P}) = \cup_{u \in U} T_{\mathbf{v}}(S_u)$. Clearly, every subcone $T_{\mathbf{v}}(S_u)$ contains a translation-invariant direction (any direction parallel to the minimal face containing $\mathbf{v}$).

Vice versa, to prove the converse implication, let us take a dissection of the tangent cone $T_{\mathbf{v}}(\mathcal{P})$ into a disjoint union of convex polytopal cones $Q_1, \dots, Q_k$. By definition of the tangent cone and since $\mathcal{P}$ can be represented as a finite union of simplices we obtain that any sufficiently small neighborhood of $\mathbf{v}$ in the tangent cone $T_{\mathbf{v}}(\mathcal{P})$ is a neighborhood of $\mathbf{v}$ in the entire $\mathcal{P}$. Consider the parallelepiped Box_{ε} centered at $\mathbf{v}$ that is ε ball centered at $\mathbf{v}$, in the L_1 -norm. Note that each convex polytopal set $Q_i \cap \text{Box}_{\varepsilon}$ can be decomposed into a union of simplices that do not contain $\mathbf{v}$ as a vertex.

Further notice that the set $\mathcal{P} \setminus \text{Box}_{\varepsilon}$ can be represented as a finite disjoint union of simplices, since Box_{ε} is the intersection of a finite number of half-spaces and $\mathcal{P}$ is a disjoint union of simplices. Clearly, every simplex in this union should not have $\mathbf{v}$ as a vertex. Now combining the dissections of each $Q_i \cap \text{Box}_{\varepsilon}$ and $\mathcal{P} \setminus \text{Box}_{\varepsilon}$ we obtain the required dissection of $\mathcal{P}$. $\square$

Proof of Proposition 6. We begin by considering the case $\rho \equiv 1$. Let $\mathcal{T}$ be a dissection of $\mathcal{P}$ with vertices $\mathcal{V}(\mathcal{T})$. Corollary 3 implies that $F_{\mathcal{P}}(\mathbf{u})$ has a denominator dividing $g_{\mathcal{T}}(\mathbf{u}) = \prod_{\mathbf{v} \in \mathcal{V}(\mathcal{T})} (1 - \langle \mathbf{v}, \mathbf{u} \rangle)$. Take $\mathbf{v}_1 \in \mathcal{V}(\mathcal{T}) \setminus \mathcal{V}(\mathcal{P})$. Then there exists another dissection $\mathcal{T}'$ such that $\mathbf{v}_1 \notin \mathcal{V}(\mathcal{T}')$. Expressing $F_{\mathcal{P}}(\mathbf{u})$ as ratios of polynomials, we have

$$F_{\mathcal{P}}(\mathbf{u}) = \frac{f_{\mathcal{T}}(\mathbf{u})}{h_{\mathcal{T}}(\mathbf{u})(1 - \langle \mathbf{v}_1, \mathbf{u} \rangle)} = \frac{f_{\mathcal{T}'}(\mathbf{u})}{g_{\mathcal{T}'}(\mathbf{u})}, \quad \text{where } g_{\mathcal{T}}(\mathbf{u}) = h_{\mathcal{T}}(\mathbf{u})(1 - \langle \mathbf{v}_1, \mathbf{u} \rangle).$$

Here $g_{\mathcal{T}'}$ is not divisible by $1 - \langle \mathbf{v}_1, \mathbf{u} \rangle$, by the choice of $\mathcal{T}'$. Thus $f_{\mathcal{T}}$ is divisible by $1 - \langle \mathbf{v}_1, \mathbf{u} \rangle$, and can be canceled out in the expression for $F_{\mathcal{P}}(\mathbf{u})$.

The case of arbitrary homogeneous ρ follows immediately by applying Theorem 1 to the already covered case $\rho \equiv 1$. $\square$

Proof of Proposition 9. First we show that for an arbitrary finite spanning set $S \subset \mathbb{R}^d$ the space $\mathfrak{M}^{\Delta}(S)$ has a basis of d -dimensional simplices containing a fixed vertex $\mathbf{v} \in S$. In particular, the set of all d -dimensional simplices containing $\mathbf{v}$ spans $\mathfrak{M}^{\Delta}(S)$ but is not necessarily a basis. Consequently, their Fantappiè transformations spans $\mathfrak{F}^{\Delta}(S)$. The following result is formulated as Theorem 4.2 of [2] and in a different form in [1]. (We omit the proof of this statement here.)

Given two points p and q and a set M in $\mathbb{R}^d$, we say that q is *visible* from p with respect to M if the straight interval pq is disjoint from M .

Lemma 15. Given a d -dimensional simplex $\sigma \subset \mathbb{R}^d$ denote by $\mathcal{V}(\sigma)$ the set of vertices of σ . Let σ^0 be the interior of σ . Let p be any point in $\mathbb{R}^d$ and let Q^+ (resp. Q^-) be the set of all $(d-1)$ -dimensional faces of σ which are visible (resp. not visible) from p with respect to σ^0 . Then the standard measures of all d -dimensional simplices with vertices in $\mathcal{V}(\sigma) \cup \{p\}$ satisfy

$$\mu_{\sigma} = \sum_{\sigma_i \in Q^+} \mu_{\sigma_i, p} - \sum_{\sigma_i \in Q^-} \mu_{\sigma_i, p},$$

where $\mu_{\sigma_i, p}$ is the standard measure of the d -dimensional simplex spanned by the vertices of σ_i and the point p .

Remark 11. If $\sigma_{i, p}$ is a degenerate simplex, i.e. p lies in the hyperplane spanned by σ_i , we simply exclude the corresponding term $\mu_{\sigma_{i, p}}$ from the above formula.

To prove Proposition 9, we need to show that $\mathfrak{F}^\Delta(S)$ coincides with $\mathfrak{Rat}(S)$ if and only if S is strongly non-degenerate. Indeed, $\mathfrak{F}^\Delta(S) \subseteq \mathfrak{Rat}(S)$ for an arbitrary spanning S , by Proposition 6. The Fantappiè transform $F_\mu : \mathfrak{M}^\Delta(S) \rightarrow \mathfrak{F}^\Delta(S)$ is a linear isomorphism which implies that $\dim \mathfrak{M}^\Delta(S) = \dim \mathfrak{F}^\Delta(S)$. By Lemma 15 the space $\mathfrak{M}^\Delta(S)$ is spanned by the standard measures μ_Δ of the set $\mathcal{B}_i$ of all d -dimensional simplices containing the fixed vertex $\mathbf{v}_i$. Let us fix the vertex $\mathbf{v}_N$ and consider the set $\mathcal{B}_N$. For S strongly non-degenerate the cardinality of $\mathcal{B}_N$ equals $\binom{N-1}{d}$.

Now we show that $\mathfrak{F}^\Delta(S) = \mathfrak{Rat}(S)$, where $\mathfrak{Rat}(S)$ has the dimension $\binom{N-1}{d}$, as it is isomorphic to the space $Pol(N-d-1, d)$ of all d -variate polynomials of degree at most $N-d-1$. This would immediately imply that the standard measures of simplices in $\mathcal{B}_N$ are linearly independent.

Lemma 16. If S is strongly non-degenerate, then $\mathfrak{F}^\Delta(S) = \mathfrak{Rat}(S)$.

Proof. We recall that $\mathfrak{F}^\Delta(S)$ comprises all linear combinations of the rational functions

$$\frac{d! \text{Vol}(\mathbf{v}_{i_1}, \dots, \mathbf{v}_{i_d}, \mathbf{v}_N)}{(1 - \langle \mathbf{v}_{i_1}, \mathbf{u} \rangle) \cdots (1 - \langle \mathbf{v}_{i_d}, \mathbf{u} \rangle) \cdot (1 - \langle \mathbf{v}_N, \mathbf{u} \rangle)}.$$

For each term $1 - \langle \mathbf{v}_i, \mathbf{u} \rangle$ we consider a (homogeneous) linear form $l_i(u_0, \mathbf{u}) = u_0 - \langle \mathbf{v}_i, \mathbf{u} \rangle$ in $d+1$ variables $u_0, \dots, u_d$. Set $n = N-1$ where $n \geq d+1$. For the n -tuple $\mathfrak{L} = \{l_1, l_2, \dots, l_n\}$ of linear $(d+1)$ -variate forms, let $V_{\mathfrak{L}}$ be the linear span of all possible products of the form $l_{i_1} l_{i_2} \dots l_{i_{n-d}}$, $1 \leq i_1 < i_2 < \dots < i_{n-d} \leq n$. Observe that $V_{\mathfrak{L}}$ is the space of all numerators that one can obtain in $\mathfrak{F}^\Delta(S)$. We need to show that $V_{\mathfrak{L}}$ contains $HPoly(n-d, d+1)$, the space of all $d+1$ -variate homogeneous polynomials of degree $n-d$. Recall that any $d+1$ -tuple of linear forms $l_{i_1}, \dots, l_{i_{d+1}}$ is linearly independent due to the strong degeneracy assumption. Thus we can express each single variable $u_0, \dots, u_d$ as a linear combination of these forms. Since $V_{\mathfrak{L}}$ contains all products $l_{i_1} l_{i_2} \dots l_{i_{n-d-1}} l_j$, where $j \in \{1, \dots, n\} \setminus \{i_1, \dots, i_{n-d-1}\}$, we conclude that $V_{\mathfrak{L}}$ contains all homogeneous polynomials of the form

$$l_{i_1} l_{i_2} \dots l_{i_{n-d-1}} u_k, \quad \text{for } 0 \leq k \leq d.$$

From that we deduce that $V_{\mathfrak{L}}$ contains all homogeneous polynomials of the form $l_{i_1} l_{i_2} \dots l_{i_{n-d-2}} u_k u_j$, where $j, k \in [n] \setminus \{i_1, \dots, i_{n-d-1}\}$. Continuing along the same lines we derive by induction that $V_{\mathfrak{L}}$ contains $HPoly(n-d, d+1)$. $\square$

For an arbitrary spanning S the cardinality of $\mathcal{B}_N$ is at most $\binom{N-1}{d} = \dim \mathfrak{Rat}(S)$. Furthermore, if S is not strongly non-degenerate the cardinality of $\mathcal{B}_N$ is strictly smaller than $\binom{N-1}{d}$, as there will be linear dependencies among the standard measures on the simplices in $\mathcal{B}_N$. Therefore $\dim \mathfrak{F}^\Delta(S) < \dim \mathfrak{Rat}(S)$. $\square$

We define the square matrix $Mat_{\mathfrak{L}}$ of size $\binom{n}{d}$ with entries being coefficients of the above products of linear forms w.r.t. the standard monomial basis in $HPol(n-d, d+1)$.

Lemma 17. The determinant of $Mat_{\mathfrak{L}}$ is proportional to the product of the determinants of all $(d+1)$ -tuples $(l_{i_1}, l_{i_2}, \dots, l_{i_{d+1}})$, $i_1 < i_2 < \dots < i_{d+1}$. (By the determinant of a $(d+1)$ -tuple of vectors in $\mathbb{R}^{d+1}$ with a fixed basis we mean the determinant of the matrix formed by the coordinates of these vectors in a chosen basis.)

Proof. Indeed, $\det(\text{Mat}_{\mathfrak{L}})$ is a form of degree $(d+1)\binom{n}{d}$ in the coefficients of the linear forms $l_1, \dots, l_n$. Thus the product $\prod_{i_1, \dots, i_{d+1}} \det(l_{i_1}, l_{i_2}, \dots, l_{i_{d+1}})$ has the same degree as $\det(\text{Mat}_{\mathfrak{L}})$. Therefore it suffices to show that $\det(\text{Mat}_{\mathfrak{L}})$ vanishes as soon as some of $\det(l_{i_1}, l_{i_2}, \dots, l_{i_{d+1}})$ vanishes. (Observe that all polynomials $\det(\text{Mat}_{\mathfrak{L}})$ are coprime.) Without loss of generality assume that l_1 is a linear combination of $l_2, \dots, l_{d+1}$. But then the column of $\text{Mat}_{\mathfrak{L}}$ corresponding to the $(n-d)$ -tuple $(1, d+2, d+3, \dots, n)$ will be a linear combination of those corresponding to $(2, d+2, d+3, \dots, n), \dots, (d+1, d+2, d+3, \dots, n)$. $\square$

Proof of Corollary 10. As we mentioned above, $\mathfrak{M}(S)$ is isomorphic to $\mathfrak{F}(S)$ and, analogously, $\mathfrak{M}^\Delta(S)$ is isomorphic to $\mathfrak{F}^\Delta(S)$. Thus, if we prove the equality $\mathfrak{F}(S) = \mathfrak{F}^\Delta(S)$ then we get $\mathfrak{M}(S) = \mathfrak{M}^\Delta(S)$. By Lemma 16, the space $\mathfrak{F}^\Delta(S)$ coincides with the linear space of all rational functions with the numerator an arbitrary polynomial of degree at most $N-d-1$ and the denominator $\Phi_S(\mathbf{u})$ equal to the product of all linear forms dual to all vertices in S . By Proposition 6 an arbitrary function in $\mathfrak{F}(S)$ is a rational function with denominator of desired form and numerator of degree at most $N-d-1$, for obvious reasons—take an arbitrary dissection and sum over its simplices. Since all such functions are already in $\mathfrak{F}^\Delta(S)$ we are done. $\square$

Proof of Theorem 12. Given a strongly non-degenerate set $S = \{\mathbf{v}_1, \dots, \mathbf{v}_{N-1}, \mathbf{v}_N\}$ and the Fantappiè transform $R(\mathbf{u}) = P(\mathbf{u})/\Phi_S(\mathbf{u})$, where $\Phi_S(\mathbf{u}) = \prod_{j=1}^N l_j(\mathbf{u})$, we want to solve the inverse moment problem. (It is easy to obtain $P(\mathbf{u})$ from information on the moments of order at most $N-d-1$ using Lemma 11.)

To solve the latter inverse problem using Corollary 3, we need to find an appropriate set of weights $\mathbf{w} = \{w_{i_1, \dots, i_d}\}$, $i_1 < i_2 < \dots < i_d$, where $w_{i_1, \dots, i_d}$ is the weight (recall Definition 1.14) of the d -dimensional simplex $\text{conv}(\mathbf{v}_{i_1}, \mathbf{v}_{i_2}, \dots, \mathbf{v}_{i_d}, \mathbf{v}_N)$ so that

$$\sum_{i_1 < i_2 < \dots < i_d} \frac{w_{i_1, \dots, i_d}}{l_{i_1}(\mathbf{u})l_{i_2}(\mathbf{u}) \dots l_{i_d}(\mathbf{u})l_N(\mathbf{u})} = \frac{P(\mathbf{u})}{l_1(\mathbf{u})l_2(\mathbf{u}) \dots l_N(\mathbf{u})}.$$

Clearing the denominators, we get the equation

$$\sum_{i_1 < i_2 < \dots < i_d} w_{i_1, \dots, i_d} l_{j_1}(\mathbf{u})l_{j_2}(\mathbf{u}) \dots l_{j_{N-d-1}}(\mathbf{u})l_N(\mathbf{u}) = P(\mathbf{u}),$$

where $\{j_1, \dots, j_{N-d-1}\} = \{1, 2, \dots, N-1\} \setminus \{i_1, \dots, i_d\}$. The latter equation is obviously equivalent to the system of linear equations

$$\text{Mat}_S \cdot \mathbf{w} = (p_{I_1}, \dots, p_{I_t}), \quad \text{where } P(\mathbf{u}) = \sum_I p_I \mathbf{u}^I,$$

and $\mathbf{w} = \{w_{i_1, \dots, i_d}\}$ is the vector consisting of the weights of all simplices containing $\mathbf{v}_N$. $\square$

Theorem 12 solves the inverse moment problem for strongly non-degenerate spanning set S . We can make this solution more explicit by giving a closed formula for the inverse matrix Mat_S^{-1} . To do this we introduce an extra variable $u_0 \in \mathbb{R}$ and identify the space $\text{Pol}(N-d-1, d)$ with the space $H\text{Pol}(N-d-1, d+1)$ of homogeneous forms of degree $N-d-1$ in $d+1$ variables $(u_0, u_1, \dots, u_d)$. We homogenize each linear form $l_i(\mathbf{u})$ in $\mathfrak{L}$ as $l_i(\mathbf{u}, u_0) = u_0 - \langle \mathbf{v}_i, \mathbf{u} \rangle$. (The matrix Mat_S remains unchanged.)

We also need the following $(d+1) \times (N-1)$ matrix $\mathbb{L}$

$$\mathbb{L} = \begin{matrix} & l_1 & l_2 & \dots & l_{N-1} \\ \begin{matrix} u_0 \\ u_1 \\ \vdots \\ u_d \end{matrix} & \begin{pmatrix} 1 & 1 & \dots & 1 \\ -\langle \mathbf{v}_1, e_1 \rangle & -\langle \mathbf{v}_2, e_1 \rangle & \dots & -\langle \mathbf{v}_{N-1}, e_1 \rangle \\ \vdots & \vdots & \dots & \vdots \\ -\langle \mathbf{v}_1, e_d \rangle & -\langle \mathbf{v}_2, e_d \rangle & \dots & -\langle \mathbf{v}_{N-1}, e_d \rangle \end{pmatrix} \end{matrix} \quad (3.1)$$

associated with Mat_S . For all possible subsets $\mathbf{i}[d] = \{i_1, \dots, i_d\}$, $i_1 < i_2 < \dots < i_d$ of d distinct columns of $\mathbb{L}$ consider the linear in $\mathbf{u}$ function $\mathbb{L}_{\mathbf{i}[d]}(\mathbf{u})$ given by:

$$\mathbb{L}_{\mathbf{i}[d]}(\mathbf{u}) = \det \begin{bmatrix} u_0 & 1 & \dots & 1 \\ u_1 & -\langle \mathbf{v}_{i_1}, e_1 \rangle & \dots & -\langle \mathbf{v}_{i_d}, e_1 \rangle \\ \vdots & \vdots & \dots & \vdots \\ u_d & -\langle \mathbf{v}_{i_1}, e_d \rangle & \dots & -\langle \mathbf{v}_{i_d}, e_d \rangle \end{bmatrix}. \quad (3.2)$$

Denote by $\mathbb{L}_{\mathbf{i}[d],j}$ the coefficient of u_j in the linear form $\mathbb{L}_{\mathbf{i}[d]}(\mathbf{u})$. For each $1 \leq j \leq N-1$ define $\mathbb{L}(j, \mathbf{i}[d])$ as

$$\mathbb{L}(j, \mathbf{i}[d]) = \det \begin{bmatrix} 1 & 1 & \dots & 1 \\ -\langle \mathbf{v}_j, e_1 \rangle & -\langle \mathbf{v}_{i_1}, e_1 \rangle & \dots & -\langle \mathbf{v}_{i_d}, e_1 \rangle \\ \vdots & \vdots & \dots & \vdots \\ -\langle \mathbf{v}_j, e_d \rangle & -\langle \mathbf{v}_{i_1}, e_d \rangle & \dots & -\langle \mathbf{v}_{i_d}, e_d \rangle \end{bmatrix}. \quad (3.3)$$

Note that if $j \notin \mathbf{i}[d]$ then $\mathbb{L}(j, \mathbf{i}[d]) \neq 0$, as by the assumption of strong non-degeneracy of S the corresponding $d+1$ linear forms are linearly independent. On the other hand, if $j \in \mathbf{i}[d]$ then we have $\mathbb{L}(j, \mathbf{i}[d]) = 0$.

The matrix Mat_S^{-1} has the following explicit description.

Lemma 18. For each $(N-d-1)$ -tuple of forms $\{l_{j_1}, l_{j_2}, \dots, l_{j_{N-d-1}}\}$ set $\mathbf{i}[d] = \{i_1, \dots, i_d\} = [N-1] \setminus \{j_1, \dots, j_{N-d-1}\}$, where $i_1 < i_2 < \dots < i_d$. Then,

$$Mat_S^{-1} = \begin{matrix} & \dots & u_0^{n_0} u_1^{n_1} \dots u_d^{n_d} & \dots \\ \vdots & & \vdots & \\ & \dots & \frac{\prod_{j=0}^d \mathbb{L}_{\mathbf{i}[d],j}^{n_j}}{\prod_{k=1}^{N-d-1} \mathbb{L}(j_k, \mathbf{i}[d])} & \dots \\ \vdots & & \vdots & \end{matrix} \quad (3.4)$$

Proof of Lemma 18. In order to show that Mat_S^{-1} defined by (3.4) is indeed the inverse of Mat_S we need to verify that $Mat_S^{-1} \cdot Mat_S$ is the identity operator on $V_{\mathcal{L}}$.

Let $\mathbf{e}'$ be the standard basis vector of $V_{\mathcal{L}}$ corresponding to the product of linear forms $l_{j'_1} \dots l_{j'_{N-d-1}}$. Then $Mat_S \cdot \mathbf{e}'$ is the vector consisting of the monomial coefficients of the homogeneous form $l_{j'_1} \dots l_{j'_{N-d-1}}$ in the variables $u_0, \dots, u_d$. Let $\mathbf{e}^T$ be the row vector of Mat_S^{-1} corresponding to the product $l_{j_1} \dots l_{j_{N-d-1}}$. We note that in $\mathbf{e}^T \cdot (Mat_S \cdot \mathbf{e}')$ one can factor out the common denominator $\prod_{k=1}^{N-d-1} \mathbb{L}(j_k, \mathbf{i}[d])$ of all fractions in $\mathbf{e}^T$; the remaining factor is of the form

$$\sum_{\substack{I=(n_0, \dots, n_d) \\ |I|=N-d-1}} Mat_S[\mathbf{u}^I, \mathbf{e}'] \prod_{j=0}^d \mathbb{L}_{\mathbf{i}[d],j}^{n_j} = l_{j'_1}(\mathbb{L}_{\mathbf{i}[d]}(\mathbf{u})) \dots l_{j'_{N-d-1}}(\mathbb{L}_{\mathbf{i}[d]}(\mathbf{u})). \quad (3.5)$$

Note that $l_{j'_k}(\mathbb{L}_{\mathbf{i}[d]}(\mathbf{u})) = \mathbb{L}(j'_k, \mathbf{i}[d])$ for $1 \leq k \leq N - d - 1$, i.e. this holds for all terms in the product on the right-hand side of (3.5). Hence, if $\mathbf{e} \neq \mathbf{e}'$ then $\mathbf{e}^T \cdot (\text{Mat}_S \cdot \mathbf{e}') = 0$, as among $j'_1, \dots, j'_{N-d-1}$ one can find $j' \in \mathbf{i}[d]$ with $\mathbb{L}(j', \mathbf{i}[d]) = 0$. On the other hand, if $\mathbf{e}$ and $\mathbf{e}'$ coincide, then the right-hand side of (3.5) is equal to $\prod_{k=1}^{N-d-1} \mathbb{L}(j_k, \mathbf{i}[d])$. Dividing by the common denominator of the fractions in $\mathbf{e}$, we obtain $\mathbf{e}^T \cdot (\text{Mat}_S \cdot \mathbf{e}) = 1$. $\square$

4. INVERSE MOMENT PROBLEM FOR WEAKLY NON-DEGENERATE S

Given an arbitrary spanning set $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_N\}$, consider the linear space $\Theta(S) \subseteq \text{Pol}(N - d - 1, d)$ spanned by all products $l_{j_1} l_{j_2} \dots l_{j_{N-d-1}}$, $j_1 < j_2 < \dots < j_{N-d-1}$. The next statement explains why we can extend our solution of the inverse moment problem from the case of strongly non-degenerate S to the case of weakly non-degenerate S .

Lemma 19. $\Theta(S) = \text{Pol}(N - d - 1, d)$ if and only if S is weakly non-degenerate, i.e. each $(d + 2)$ -tuple of points of S is spanning.

Proof. We have N (non-homogeneous) linear forms $l_1, \dots, l_N$ in variables $\mathbf{u} = (u_1, \dots, u_d)$ and the linear space $V_{\mathcal{L}}$ spanned by all possible products of $(N - d - 1)$ -tuples of distinct forms. We need to investigate whether $V_{\mathcal{L}}$ coincides with $\text{Pol}(N - d - 1, d)$. Homogenizing, we consider the same question for the linear homogeneous forms and the homogeneous polynomials of degree $N - d - 1$ in variables $(u_0, u_1, \dots, u_d)$.

First assume that there are $d + 2$ linear forms $l_1, \dots, l_{d+2}$ which are not spanning. Then one can find a non-zero vector $\mathbf{z}_0 \in \mathbb{R}^{d+1}$, such that $l_1(z) = \dots = l_{d+2}(z) = 0$. Note that each product of $N - d - 1$ different forms chosen from $l_1, \dots, l_N$ contains at least one form among $\{l_1, \dots, l_{d+2}\}$. Therefore any linear combination of products of $N - d - 1$ forms vanishes at $\mathbf{z}_0$. Thus $V_{\mathcal{L}}$ cannot coincide with $H\text{Pol}(N - d - 1, d + 1)$.

Conversely, assume that every $(d + 2)$ -tuple of distinct forms among $l_1, \dots, l_N$ is spanning. First, we notice that $H\text{Pol}(N - d - 1, d + 1)$ can be spanned by the all possible products of $N - d - 1$ linear forms (not necessarily pairwise distinct). Indeed, since first $d + 2$ forms span the dual space of $\mathbb{R}^{d+1}$, we can express each variable x_i as a linear combination of these forms. Therefore every monomial of degree $N - d - 1$ can be expressed as a linear combination of products of $N - d - 1$ forms.

Now we show that each product of $N - d - 1$, not necessarily distinct, forms can be expressed as a linear combination of the products of distinct ones. Assume the contrary and consider monomials $l_1^{i_1} \dots l_N^{i_N}$ of degree $N - d - 1$ which cannot be expressed as a linear combination of products with all distinct forms. Among those monomials we take a monomial $\mathbf{m} = l_1^{k_1} \dots l_N^{k_N}$ having the maximal number of distinct forms in the product. Since $\mathbf{m}$ is not a product of all distinct forms, it should contain a form l_i in some power $k_i \geq 2$. Given that $k_i \geq 2$ and the degree of $\mathbf{m}$ is $N - d - 1$, one can find $d + 2$ distinct forms $l_{i_1}, \dots, l_{i_{d+2}}$ that do not appear in $\mathbf{m}$. Since any $d + 2$ of our forms span the dual space of $\mathbb{R}^{d+1}$, we can express l_i as a linear combination of $l_{i_1}, \dots, l_{i_{d+2}}$. Now rewrite $\mathbf{m}$ as $(\alpha_1 \cdot l_{i_1} + \dots + \alpha_{d+2} \cdot l_{i_{d+2}}) l_1^{k_1} \dots l_i^{k_i-1} \dots l_N^{k_N}$, where $\alpha_1 \cdot l_{i_1} + \dots + \alpha_{d+2} \cdot l_{i_{d+2}} = l_i$. Thus we get an expression of $\mathbf{m}$ as a linear combination of monomials $\alpha_j \cdot l_{i_j} l_1^{k_1} \dots l_i^{k_i-1} \dots l_N^{k_N}$, where each such monomial has more distinct forms than $\mathbf{m}$. Each of such monomials can be expressed as a linear combination of products of all distinct forms, since $\mathbf{m}$ was chosen as a monomial with the maximal possible

number of distinct forms, which cannot be expressed in such a way. This is a contradiction. Therefore $\mathbf{m}$ can also be expressed as a linear combination of products of all distinct forms. $\square$

Below we consider the inverse problem for a weakly non-degenerate S , using notation from (3.1) and (3.3). Here we no longer have a natural basis of all simplices sharing a common vertex $\mathbf{v}_N$. Because of that we need to consider all N points and include one more linear form l_N into the corresponding matrix $\mathbb{L}$. Slightly abusing our notation, we denote by $\mathbb{L}$ the same matrix as before, although it contains one more (last) column corresponding to $\mathbf{v}_N$. Similarly to notation (3.3), for a given set J of $d+1$ linear forms we denote by $\mathbb{L}(J)$ the determinant of the corresponding $(d+1) \times (d+1)$ -minor of $\mathbb{L}$.

We introduce the extended $\binom{N-1}{d} \times \binom{N}{d+1}$ -matrix $\widetilde{Mat}_S$ with columns consisting of the coefficients of the homogeneous polynomial $l_{i_1}(\mathbf{u}) \dots l_{i_{N-d-1}}(\mathbf{u})$ with respect to the monomial basis in the variables $(u_0, u_1, \dots, u_d)$. By Lemma 19, $\widetilde{Mat}_S$ has full rank, since it determines a surjective linear map *onto* $H\text{Pol}(N-d-1, d+1)$. Thus $\widetilde{Mat}_S$ has a maximal minor with a non-vanishing determinant. Formula (4.1) holds for the determinant of any maximal minor of $\widetilde{Mat}_S$.

Lemma 20. Let $\mathfrak{S}$ be any set of $\binom{N-1}{d}$ columns of $\widetilde{Mat}_S$. We label each column $T \in \mathfrak{S}$ by the corresponding subset of the linear forms $l_1, \dots, l_N$ of cardinality $N-d-1$. Then the determinant of the maximal minor $\widetilde{Mat}_S(\mathfrak{S})$ formed by the columns of $\mathfrak{S}$ is given by:

$$\det [\widetilde{Mat}_S(\mathfrak{S})] = k(\mathfrak{S}) \cdot \prod_{\substack{J \in \binom{N}{d+1}: \\ \forall T \in \mathfrak{S} \ T \cap J \neq \emptyset}} \mathbb{L}(J), \quad (4.1)$$

where $k(\mathfrak{S})$ is a constant (possibly equal to zero) depending only on the combinatorial structure of the $(N-d-1)$ -tuples in the set $\mathfrak{S}$.

Proof. Fix the set $\mathfrak{S}$ as above. In what follows we treat both sides of (4.1) as complex-valued polynomials in $N \cdot (d+1)$ variables, these variables being the entries of matrix $\mathbb{L}$.

We first show that every determinant $\mathbb{L}(J)$ divides $\det [\widetilde{Mat}_S(\mathfrak{S})]$. Indeed, let $J = \{j_1, \dots, j_{d+1}\}$ be a set of $(d+1)$ forms which has a nonempty intersection with any $(N-d-1)$ -tuple of forms in $\mathfrak{S}$. Let $\mathbf{z} = (z_1, \dots, z_{N \cdot (d+1)})$ be a zero of the polynomial $\mathbb{L}(J)$, which means that forms $l_{j_1}, \dots, l_{j_{d+1}}$ comprised of the corresponding coordinates of $\mathbf{z}$ are linearly dependent. Therefore, there is a non-zero vector $\mathbf{u}_0 \in \mathbb{R}^{d+1}$, such that $l_{j_1}(\mathbf{u}_0) = \dots = l_{j_{d+1}}(\mathbf{u}_0) = 0$. Consider the row vector $(\mathbf{u}_0^T)$ consisting of $\binom{N-1}{d}$ homogeneous monomials of degree $N-d-1$ evaluated at $\mathbf{u}_0$. We notice that $(\mathbf{u}_0^T)$ is in the kernel of $\widetilde{Mat}_S(\mathfrak{S})$, as the product of $(\mathbf{u}_0^T)$ with each column vector $T \in \mathfrak{S}$ of $\widetilde{Mat}_S(\mathfrak{S})$ is equal to $\prod_{j \in T} l_j(\mathbf{u}_0)$; and every set $T \in \mathfrak{S}$ contains at least one of the forms $l_{j_1}, \dots, l_{j_{d+1}}$ in such a product. Thus $\det [\widetilde{Mat}_S(\mathfrak{S})]$ also vanishes at such $\mathbf{z}$.

We recall a well-known fact (cf. e.g. [10, Theorem 61.1]) that

$$\mathbb{L}(J) = \det \begin{pmatrix} & l_{j_1} & l_{j_2} & \dots & l_{j_{d+1}} \\ u_0 & z_{j_1} & z_{j_2} & \dots & z_{j_{d+1}} \\ u_1 & z_{j_1+N} & z_{j_2+N} & \dots & z_{j_{d+1}+N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ u_d & z_{j_1+dN} & z_{j_2+dN} & \dots & z_{j_{d+1}+dN} \end{pmatrix},$$

is an irreducible complex-valued polynomial in variables $z_{j_1}, z_{j_2}, \dots, z_{j_{d+1}+Nd}$. Now if every zero of an irreducible polynomial $p(z_1, \dots, z_{N(d+1)})$ annihilates another polynomial $q(z_1, \dots, z_{N(d+1)})$ then p divides q . We conclude that $\mathbb{L}(J)$ divides $\det [\widetilde{Mat}_S(\mathfrak{S})]$.

Using the fact that each $\mathbb{L}(J)$ is an irreducible polynomial and all $\mathbb{L}(J)$'s are pairwise distinct (i.e. have distinct sets of projective zeros) we conclude that the product of $\mathbb{L}(J)$'s in the right-hand side of (4.1) divides $\det [\widetilde{Mat}_S(\mathfrak{S})]$.

Finally, the product of $\mathbb{L}(J)$'s has the degree

$$(d+1) \left| \left\{ J \in \left[\begin{matrix} N \\ d+1 \end{matrix} \right] \mid \forall T \in \mathfrak{S} \quad T \cap J \neq \emptyset \right\} \right|. \quad (4.2)$$

We observe that for each $T \in \mathfrak{S}$ the complementary set of $d+1$ forms cannot be taken as a feasible J . We notice further that these complements are the only exceptions for the choice of J . Therefore as a feasible J we can pick any of $\binom{N}{d+1}$ $(d+1)$ -tuples except those $\binom{N-1}{d}$ complements of a $T \in \mathfrak{S}$. Therefore, (4.2) equals

$$(d+1) \left(\binom{N}{d+1} - \binom{N-1}{d} \right) = (d+1) \binom{N-1}{d+1} = (N-d-1) \binom{N-1}{d}.$$

The latter expression coincides with the degree of the polynomial $\det [\widetilde{Mat}_S(\mathfrak{S})]$ (assuming that it is not a zero), as $\widetilde{Mat}_S(\mathfrak{S})$ has $\binom{N-1}{d}$ columns and each entry is a homogeneous polynomial of degree $N-d-1$.

Hence $\widetilde{Mat}_S(\mathfrak{S})$ coincides with the product of $\mathbb{L}(J)$'s up to a constant factor which might vanish. This constant does not depend on the entries of matrix $\mathbb{L}$ and hence it is completely determined by the set $\mathfrak{S}$, regardless of the location of points of S in $\mathbb{R}^d$. $\square$

Lemma 19 allows us to solve the inverse moment problem for a given weakly non-degenerate $S = \{\mathbf{v}_1, \dots, \mathbf{v}_N\}$ in a certain linear space $\widetilde{\mathfrak{M}}(S) \supseteq \mathfrak{M}^\Delta(S)$ of measures supported on $\text{conv}(S)$. Namely, $\widetilde{\mathfrak{M}}(S)$ is spanned by measures $\mu \in \widetilde{\mathfrak{M}}(S)$ whose normalized moment generating functions $F_\mu(\mathbf{u})$ belong to $\mathfrak{Rat}(S)$, i.e. $F_\mu(\mathbf{u}) = P(\mathbf{u})/\Phi_S(\mathbf{u})$, where $\Phi_S(\mathbf{u}) = \prod_{j=1}^N l_j(\mathbf{u})$ and $P(\mathbf{u})$ is a polynomial of degree at most $N-d-1$. Indeed, by Lemma 19 any $R(\mathbf{u}) \in \mathfrak{Rat}(S)$ can be represented in the form

$$R(\mathbf{u}) = \sum_{i_1 < i_2 < \dots < i_{d+1} \leq N} \frac{K_{i_1 i_2 \dots i_{d+1}}}{l_{i_1} l_{i_2} \dots l_{i_{d+1}}}, \quad (4.3)$$

with some real constants $K_{i_1 i_2 \dots i_{d+1}}$. If $\mathbf{v}_{i_1}, \mathbf{v}_{i_2}, \dots, \mathbf{v}_{i_{d+1}}$ span $\mathbb{R}^d$ then the term $\frac{K_{i_1 i_2 \dots i_{d+1}}}{l_{i_1} l_{i_2} \dots l_{i_{d+1}}}$ can be interpreted as the normalized moment generating function of an appropriately scaled standard measure of the d -dimensional simplex spanned by these vertices.

If $\mathbf{v}_{i_1}, \mathbf{v}_{i_2}, \dots, \mathbf{v}_{i_{d+1}}$ only span a hyperplane H in $\mathbb{R}^d$ then (4.3) corresponds to a singular (w.r.t. to the Lebesgue measure on $\mathbb{R}^d$) measure μ_δ supported on $\delta = \text{conv}(\mathbf{v}_{i_1}, \dots, \mathbf{v}_{i_{d+1}})$. One way to define it as the weak limit of a sequence of (absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}^d$) measures—the appropriately scaled standard measures μ_{δ_t} of family of d -dimensional simlices δ_t which degenerate into δ when $t = 0$. There is no loss in generality in assuming $K_{i_1 i_2 \dots i_{d+1}} = 1$, i.e. to deal with probability measures.

Proposition 21. Let $\mathbf{W} = \{\mathbf{w}_1, \dots, \mathbf{w}_d, \mathbf{w}_{d+1}\}$ be a $(d+1)$ -tuple of points in $\mathbb{R}^d$ such that $\mathbf{W}$ spans a hyperplane $H \subset \mathbb{R}^d$. Denote by $l_{\mathbf{w}_1} = 1 - \langle \mathbf{w}_1, \mathbf{u} \rangle, \dots, l_{\mathbf{w}_{d+1}} =$

$1 - \langle \mathbf{w}_{d+1}, \mathbf{u} \rangle$ the associated linear forms. There exists a unique measure $\mu_{\mathbf{W}}$ supported on $\delta = \text{conv}(\mathbf{W})$ with the normalized moment generating function $F_{\mu_{\mathbf{W}}}(\mathbf{u})$ given by

$$F_{\mu_{\mathbf{W}}}(\mathbf{u}) = \frac{1}{l_{\mathbf{w}_1} l_{\mathbf{w}_2} \dots l_{\mathbf{w}_{d+1}}}. \quad (4.4)$$

Proof. Without loss of generality assume that $\mathbf{W} = \{\mathbf{w}_1, \dots, \mathbf{w}_d, \mathbf{w}_{d+1}\}$ is ordered in such a way that $\{\mathbf{w}_1, \dots, \mathbf{w}_d\}$ span H . Then, δ_t is defined as $\delta_t = \text{conv}(\delta, \mathbf{w}_{d+1} + t\mathbf{z})$, with $\mathbf{z}$ a unit normal to H , and μ_{δ_t} as the uniform density probability measure supported on δ_t . Then $\lim_{t \rightarrow 0} \mu_{\delta_t} = \mu_{\delta}$, where $\lim$ is understood in sense of *weak convergence of distributions (measures)*, i.e. that $\lim_{t \rightarrow 0} \int f d\mu_{\delta_t} = \int f d\mu_{\delta}$ for any bounded, continuous real function on $\mathbb{R}^d$, cf. e.g. [9]. Then, this measure has compact support, and thus is determined by its moments, cf. e.g. [24, Proposition 3.2]. $\square$

Remark 12. One can prove that the integration of a smooth compactly supported function ϕ with respect to the limiting measure $\mu_{\mathbf{W}}$ is given by the integration of ϕ over δ with a continuous piecewise linear weight function uniquely determined by δ . Similar limits appear frequently in the theory of splines. Since we only need the existence of $\mu_{\mathbf{W}}$ we do not pursue this topic here.

Our solution of the inverse moment problem for the linear space $\widetilde{\mathfrak{M}}(S)$ closely follows the pattern presented in Example 2. In other words, given a weakly non-degenerate S and the set of moments up to order $N - d - 1$ we

- (i) produce the rational function $R(\mathbf{u}) \in \mathfrak{Rat}(S)$ with Taylor coefficients coinciding with the normalized moments;
- (ii) represent $R(\mathbf{u})$ in the form (4.3);
- (iii) for each term as in (4.4) determine the underlying measure supported on the (probably degenerate) convex hull of the vertices $\mathbf{v}_{i_1}, \mathbf{v}_{i_2}, \dots, \mathbf{v}_{i_{d+1}}$.

We can now prove our central result claiming that $\mathfrak{M}^{\Delta}(S) = \mathfrak{M}(S)$ for a weakly non-degenerate S .

Proof of Theorem 8. Theorem 8 is already settled in Corollary 10 for the case of strongly non-degenerate S . It remains to consider the case of weakly non-degenerate S . The denominator of the moment generating function $F_{\mathcal{P}}(\mathbf{u})$ for an arbitrary generalized polytope $\mathcal{P}$ with the vertex set S is of the form $\prod_{i=1}^N l_i$ by Proposition 6, and its numerator belongs to $\text{Pol}(N - d - 1, d)$. As S is weakly non-degenerate, $F_{\mathcal{P}}(\mathbf{u})$ can be written as a linear combination of the fractions as in (4.3), where $(i_1, i_2, \dots, i_{d+1})$ runs over the set of $(d + 1)$ -tuples of indices. If a $(d + 1)$ -tuple $l_{i_1}, l_{i_2}, \dots, l_{i_{d+1}}$ is spanning then $\frac{K}{l_{i_1} l_{i_2} \dots l_{i_{d+1}}}$ is the moment generating function of the measure supported on the simplex Δ , determined by its denominator, with the uniform density $K/d! \text{Vol}(\Delta)$. By Proposition 21, if a $(d + 1)$ -tuple $l_{i_1}, l_{i_2}, \dots, l_{i_{d+1}}$ is not spanning then $\frac{K}{l_{i_1} l_{i_2} \dots l_{i_{d+1}}}$ is the moment generating function of a singular measure supported on a degenerate simplex. As $\mathcal{P}$ is a generalized polytope, its standard measure has no singular components. Therefore, no degenerate simplices can appear in its decomposition. $\square$

Remark 13. The latter proof demonstrates that if one starts from the set of moments of the standard measure μ of a polytope with the vertex set S then we never obtain degenerate simplices while solving the inverse moment problem. This is why $\mathfrak{M}^{\Delta}(S) = \mathfrak{M}(S)$. However, an explicit description of $\mathfrak{F}^{\Delta}(S)$ for a general weakly non-degenerate S is missing at present. For concrete Examples 3 and 4 we give these descriptions below.

Our final result computes $\dim \mathfrak{M}^\Delta(S)$ and describes a procedure to construct a basis for $\mathfrak{M}^\Delta(S)$.

Proposition 22. Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_N\} \subset \mathbb{R}^d$ be an arbitrary weakly non-degenerate spanning set. Then

- (i) $\dim \mathfrak{M}^\Delta(S) = \binom{N-1}{d} - \sharp_{deg}$ where $\sharp_{deg}$ is the number of degenerate simplices, i.e. the number of non-spanning $(d+1)$ -tuples of points of S .
- (ii) If δ is a degenerate d -dimensional simplex with vertices in $S \setminus \{\mathbf{v}_i\}$ then there is exactly one linear dependence among the standard measures of all d -dimensional simplices on $\mathbf{v}_i$ and d vertices of δ .
- (iii) The standard measure of any d -dimensional simplex on $\mathbf{v}_i$ is contained in at most one dependence as in (ii).
- (iv) For any vertex $\mathbf{v}_i$ one can construct a (in general, non-unique) basis $\mathcal{B}_i$ of $\mathfrak{M}^\Delta(S)$ consisting of standard measures of d -dimensional simplices on $\mathbf{v}_i$, as follows.
 - (a) Start from the set $\mathcal{B}_i$ of the d -dimensional simplices on $\mathbf{v}_i$.
 - (b) For each degenerate simplex δ not containing $\mathbf{v}_i$, remove from $\mathcal{B}_i$ the standard measure of an arbitrary simplex on $\mathbf{v}_i$ from the corresponding to δ linear dependence, cf. (i).

Thus we obtain $\binom{N-1}{d} - \sharp_{deg}$ standard measures of d -dimensional simplices, forming a basis of $\mathfrak{M}^\Delta(S)$.

Proof. To prove (i), notice that $\dim \widetilde{\mathfrak{M}}(S) = \dim \mathfrak{Rat}(S) = \binom{N-1}{d}$. As well, $\widetilde{\mathfrak{M}}(S) = \mathfrak{M}(S) \oplus \mathfrak{M}_{deg}(S)$, where $\mathfrak{M}_{deg}(S)$ is the linear span of the measures $\mu_\delta^{(1)}$ with δ running over the set of all degenerate simplices spanned by $(d+1)$ -tuples of dependent vertices in S , cf. Proposition 21. Observe that these measures $\mu_\delta^{(1)}$ are linearly independent, as each degenerate simplex defines a singular measure supported in a proper hyperplane, and these hyperplanes differ for different degenerate simplices. We are done with (i).

Let Σ_0 be a dependent $d+1$ -subset of S , and $\delta = \text{conv}(\Sigma_0)$ be as in (ii). Then δ spans a hyperplane H_0 . As each d -dimensional simplex on $\sigma_0 := \mathbf{v}_i$ and d vertices from Σ_0 is uniquely defined by the latter, it suffices to analyze dependencies between the standard measures of $d-1$ -simplices with vertices in Σ_0 .

We can view Σ_0 as a weakly non-degenerate subset in $\mathbb{R}^{d-1} \cong H_0$. By (i), we have $\dim \mathfrak{M}^\Delta(\Sigma_0) = \binom{d}{d-1} - \sharp_{deg}(\Sigma_0)$. If Σ_0 is strongly non-degenerate as a subset of $H_0 \cong \mathbb{R}^{d-1}$, i.e. $\sharp_{deg}(\Sigma_0) = 0$, then $\dim \mathfrak{M}^\Delta(\Sigma_0) = d$, i.e. there is exactly one linear dependence between the standard measures of $d-1$ -simplices with vertices in Σ_0 , and we are done. Otherwise, $\Sigma_0 = \{\sigma_1\} \cup \Sigma_1$, with Σ_1 spanning a hyperplane H_1 in H_0 . Moreover, this can only happen if $d \geq 3$. Now, we can repeat the whole argument with σ_1 in place of σ_0 , Σ_1 in place of Σ_0 , and H_1 in place of H_0 . Again, we either have Σ_1 strongly degenerate, and we are done, or we repeat this argument, etc., until we hit a strongly non-degenerate Σ_k , which is bound to happen, as the dimension goes down each iteration. This completes the proof of (ii).

Then, (iii) stems from the fact that the vertices of d -dimensional simplex on $\mathbf{v}_i$ distinct from $\mathbf{v}_i$ span a hyperplane, and the only possibility for a degenerate simplex δ as in (ii) is to lie in this hyperplane.

Finally, to prove (iv), observe that the set $\mathcal{B}'_i$ of the standard measures of d -dimensional simplices containing a given vertex $\mathbf{v}_i$ always spans $\mathfrak{M}^\Delta(S)$, see Lemma 15. Now for each degenerate d -simplex δ we prune $\mathcal{B}'_i$ by removing the standard measure of a simplex in the linear dependence corresponding to δ . In view of (ii) and (iii) this process is well-defined and unambiguous. In the end we obtain $\binom{N-1}{d} - \sharp_{deg}$ standard measures of d -dimensional simplices. In view of (i) they form a basis of $\mathfrak{M}^\Delta(S)$, as claimed. $\square$

Remark 14. The above discussions show that the columns of $\widetilde{Mat}_S$ corresponding to degenerate simplices must necessarily be included in any non-vanishing maximal minor $\widetilde{Mat}_S(\mathfrak{S})$.

We conclude our discussion of the weakly non-degenerate case with two examples.




FIGURE 3. Vertices for Examples 3 and 4.

Example 3. Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4, \mathbf{v}_5\}$ where $\mathbf{v}_1 = (1, 1), \mathbf{v}_2 = (2, 0), \mathbf{v}_3 = (2, 2), \mathbf{v}_4 = (0, 2), \mathbf{v}_5 = (0, 0)$. Then $l_1 = 1 - u_1 - u_2, l_2 = 1 - 2u_1, l_3 = 1 - 2u_1 - 2u_2, l_4 = 1 - 2u_2, l_5 = 1$. Calculating the products $l_i l_j, i < j$ and taking their coefficients in the standard monomial basis of $Pol(2, 2)$ we obtain the following 6×10 -matrix $\widetilde{Mat}_S$

$$\widetilde{Mat}_S = \begin{matrix} & l_1 l_2 & l_1 l_3 & l_1 l_4 & l_1 l_5 & l_2 l_3 & l_2 l_4 & l_2 l_5 & l_3 l_4 & l_3 l_5 & l_4 l_5 \\ \begin{matrix} 1 \\ u_1 \\ u_2 \\ u_1^2 \\ u_1 u_2 \\ u_2^2 \end{matrix} & \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ -3 & -3 & -1 & -1 & -4 & -2 & -2 & -2 & -2 & 0 \\ -1 & -3 & -3 & -1 & -2 & -2 & 0 & -4 & -2 & -2 \\ 2 & 2 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 0 \\ 2 & 4 & 2 & 0 & 4 & 4 & 0 & 4 & 0 & 0 \\ 0 & 2 & 2 & 0 & 0 & 0 & 0 & 4 & 0 & 0 \end{pmatrix} \end{matrix}.$$

Its rank equals 6 and one of non-vanishing maximal minors consists of the columns with numbers $\mathfrak{S} = \{5, 6, 7, 8, 9, 10\}$. (Recall that any non-vanishing maximal minor must include columns 6 and 9 corresponding to degenerate triples $(\mathbf{v}_1, \mathbf{v}_3, \mathbf{v}_5)$ and $(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_4)$ resp.) The corresponding submatrix $\widetilde{Mat}_S(\mathfrak{S})$ equals

$$\widetilde{Mat}_S(\mathfrak{S}) = \begin{matrix} & l_2 l_3 & l_2 l_4 & l_2 l_5 & l_3 l_4 & l_3 l_5 & l_4 l_5 \\ \begin{matrix} 1 \\ u_1 \\ u_2 \\ u_1^2 \\ u_1 u_2 \\ u_2^2 \end{matrix} & \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ -4 & -2 & -2 & -2 & -2 & 0 \\ -2 & -2 & 0 & -4 & -2 & -2 \\ 4 & 0 & 0 & 0 & 0 & 0 \\ 4 & 4 & 0 & 4 & 0 & 0 \\ 0 & 0 & 0 & 4 & 0 & 0 \end{pmatrix} \end{matrix}.$$

Further,

$$4\widetilde{Mat}_S^{-1}(\mathfrak{S}) = \begin{matrix} & 1 & u_1 & u_2 & u_1^2 & u_1u_2 & u_2^2 \\ \begin{matrix} l_2l_3 \\ l_2l_4 \\ l_2l_5 \\ l_3l_4 \\ l_3l_5 \\ l_4l_5 \end{matrix} & \begin{pmatrix} 0 & 0 & 4 & 0 & -4 & 4 \\ 0 & 0 & 0 & 0 & -2 & 2 \\ 0 & 0 & 2 & 0 & -2 & 0 \\ 1 & -1 & 0 & 0 & -1 & 1 \\ 0 & 1 & 0 & 0 & -1 & 0 \\ 0 & -1 & 1 & 1 & -1 & 0 \end{pmatrix} \end{matrix}.$$

Thus, given an arbitrary rational function $R(u_1, u_2) = P(u_1, u_2)/\Phi_S(u_1, u_2)$ where $P(u_1, u_2) = a_{00} + a_{1,0}u_1 + a_{0,1}u_2 + a_{2,0}u_1^2 + a_{11}u_1u_2 + a_{02}u_2^2$ is any polynomial of degree at most 2 and $\Phi_S(u_1, u_2) = l_1l_2l_3l_4l_5$, we obtain

$$\begin{cases} w_{145} = -a_{10} - a_{20} - a_{11} \\ w_{135} = -\frac{1}{2}(a_{11} - a_{02}) \\ w_{134} = \frac{1}{2}(a_{01} - a_{11}) \\ w_{125} = \frac{1}{4}(a_{00} - a_{10} - a_{11} + a_{02}) \\ w_{124} = \frac{1}{4}(a_{10} - a_{11}) \\ w_{123} = -\frac{1}{4}(a_{10} - a_{01} - a_{20} + a_{11}). \end{cases}$$

Triangles Δ_{135} and Δ_{124} are degenerate which implies that if the original measure we are recovering is polygonal then $w_{135} = w_{124} = 0$. Therefore the linear space of numerators $P(u_1, u_2)$ for the space $\mathfrak{F}^\Delta(S)$ in this example is given by the relation

$$a_{01} = a_{11} = a_{02}.$$

Our last example is more degenerate than the previous one, although still weakly non-degenerate. In fact, in this example S is a multiset since $\mathbf{v}_1 = \mathbf{v}_5$. It shows that our technique can be generalized to a certain class of multisets as well.

Example 4. Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4, \mathbf{v}_5\}$ where $\mathbf{v}_1 = \mathbf{v}_5 = (0, 0)$, $\mathbf{v}_2 = (2, 0)$, $\mathbf{v}_3 = (1, 1)$, $\mathbf{v}_4 = (0, 2)$. Then $l_1 = l_5 = 1$, $l_2 = 1 - 2u_1$, $l_3 = 1 - u_1 - u_2$, $l_4 = 1 - 2u_2$. Calculating all products $l_i l_j$, $i < j$ and taking their coefficients in the standard monomial basis of $Pol(2, 2)$, we obtain the following 6×10 -matrix $\widetilde{Mat}_S$:

$$\widetilde{Mat}_S = \begin{matrix} & l_1l_2 & l_1l_3 & l_1l_4 & l_1l_5 & l_2l_3 & l_2l_4 & l_2l_5 & l_3l_4 & l_3l_5 & l_4l_5 \\ \begin{matrix} 1 \\ u_1 \\ u_2 \\ u_1^2 \\ u_1u_2 \\ u_2^2 \end{matrix} & \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ -2 & -1 & 0 & 0 & -3 & -2 & 0 & -1 & -1 & 0 \\ 0 & -1 & -2 & 0 & -1 & -2 & -2 & -3 & -1 & -2 \\ 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 4 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 0 & 0 \end{pmatrix} \end{matrix}.$$

Its rank equals 6 and a non-vanishing maximal minor consists of the columns with numbers $\mathfrak{S} = \{1, 3, 4, 5, 6, 8\}$. The corresponding submatrix $\widetilde{Mat}_S(\mathfrak{S})$ equals

$$\widetilde{Mat}_S(\mathfrak{S}) = \begin{matrix} & l_1l_2 & l_1l_4 & l_1l_5 & l_2l_3 & l_2l_4 & l_3l_4 \\ \begin{matrix} 1 \\ u_1 \\ u_2 \\ u_1^2 \\ u_1u_2 \\ u_2^2 \end{matrix} & \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ -2 & 0 & 0 & -3 & -2 & -1 \\ 0 & -2 & 0 & -1 & -2 & -3 \\ 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 2 & 4 & 2 \\ 0 & 0 & 0 & 0 & 0 & 2 \end{pmatrix} \end{matrix}.$$

Further,

$$4\widetilde{Mat}_S^{-1}(\mathfrak{S}) = \begin{matrix} & 1 & u_1 & u_2 & u_1^2 & u_1u_2 & u_2^2 \\ \begin{matrix} l_1l_2 \\ l_1l_4 \\ l_1l_5 \\ l_2l_3 \\ l_2l_4 \\ l_3l_4 \end{matrix} & \begin{pmatrix} 0 & -2 & 0 & -2 & -1 & 0 \\ 0 & 0 & -2 & 0 & -1 & -2 \\ 4 & 2 & 2 & 1 & 1 & 1 \\ 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 2 \end{pmatrix} \end{matrix}.$$

Thus, given an arbitrary rational function $R(u_1, u_2) = P(u_1, u_2)/\Phi_S(u_1, u_2)$ where $P(u_1, u_2) = a_{00} + a_{1,0}u_1 + a_{0,1}u_2 + a_{2,0}u_1^2 + a_{11}u_1u_2 + a_{02}u_2^2$ is any polynomial of degree at most 2 and $\Phi_S(u_1, u_2) = l_1l_2l_3l_4l_5$, we obtain

$$\begin{cases} w_{345} = -\frac{1}{4}(2a_{10} + 2a_{20} + a_{11}) \\ w_{235} = -\frac{1}{4}((2a_{01} + a_{11} + 2a_{02}) \\ w_{234} = 4a_{00} + 2a_{01} + 2a_{10} + a_{20} + a_{11} + a_{02} \\ w_{145} = 2a_{20} \\ w_{135} = -a_{10} + a_{11} - a_{20} \\ w_{125} = 2a_{02}. \end{cases}$$

Notice that triangles $\Delta_{125}, \Delta_{135}, \Delta_{145}, \Delta_{234}$ are degenerate. If we know that the original measure we are recovering is polygonal then one should get $w_{125} = w_{135} = w_{145} = w_{234} = 0$. Therefore, the linear space of numerators $P(u_1, u_2)$ for the space $\mathfrak{F}^\Delta(S)$ in this example is given by the system of equations:

$$\begin{cases} a_{20} = 0 \\ a_{10} = a_{11} \\ a_{02} = 0 \\ 4a_{00} + 2a_{01} + 3a_{10} = 0. \end{cases}$$

5. REMARKS AND OPEN PROBLEMS

Remark 15. A weaker form of Corollary 4 (i.e. the rationality of $F_{\mathcal{P}}^p(\mathbf{u})$, but without the claim on the particular shape of the denominator) can be derived directly from (1.3) by using Stokes formula, along the lines of [6, Lemma 1].

Problem 2. Find an appropriate version of Theorem 2, applicable to non-simple and/or non-convex polytopes.

Remark 16. Choose an arbitrary basis $\{\Delta_j\}$ of $\mathfrak{M}^\Delta(S)$ consisting of the standard measures of simplices. The set $\{\Delta_j\}$ spans an integer lattice in $\mathfrak{M}^\Delta(S)$. (One can easily see that this lattice is invariantly defined independently of the choice of a basis of standard measures of simplices.) Denote by $\mathfrak{M}_{\mathbb{Z}}^\Delta(S)$ the space $\mathfrak{M}^\Delta(S)$ with the latter lattice. We can prove the following.

Proposition 23. If Conjecture 7 holds then any generalized polytope $\mathcal{P} \in \mathcal{P}(S)$ with standard measure $\mu_{\mathcal{P}}$ in $\mathfrak{M}^\Delta(S)$ corresponds to a rational point in $\mathfrak{M}_{\mathbb{Z}}^\Delta(S)$.

Proof. One can easily show that any $\mathcal{P}$ as above can be represented as the union of the closures of connected components of $\mathbb{R}^d \setminus H(S)$, where $H(S)$ is the hyperplane arrangement consisting of all hyperplanes spanned by d -tuples of points in S . (The converse is obviously not true.) Let $\tilde{S} \supseteq S$ be the *extended set of vertices* obtained by adding to S all vertices of the hyperplane arrangement $H(S)$. Since each connected component in $\mathbb{R}^d \setminus H(S)$ is convex, it can be triangulated on $\tilde{S}$. Consider the space $\mathfrak{M}_{\mathbb{Z}}^\Delta(\tilde{S})$. Obviously, $\mu_{\mathcal{P}}$ is an integer point in $\mathfrak{M}_{\mathbb{Z}}^\Delta(\tilde{S})$. Also, $\mathfrak{M}_{\mathbb{Z}}^\Delta(S)$ is

contained in $\mathfrak{M}_{\mathbb{Z}}^{\Delta}(\tilde{S})$ as a sublattice. Thus, if $\mu_{\mathcal{P}}$ belongs to $\mathfrak{M}_{\mathbb{Z}}^{\Delta}(S)$ it is a rational point there. $\square$

Remark 17. If either $\mathcal{P}$ itself or its complement in $\text{conv}(S)$ can be triangulated by simplices with vertices in S then $\mu_{\mathcal{P}}$ is an integer point in $\mathfrak{M}_{\mathbb{Z}}(S)$.

Let us pose the following tantalizing question.

Problem 3. Does there exist a generalized polytope $\mathcal{P}$ with the set of vertices $S(\mathcal{P})$ and with non-integer coordinates in $\mathfrak{M}_{\mathbb{Z}}(S(\mathcal{P}))$?

Problem 4. One can also define a rational convex cone $\mathfrak{Pos}(S) \subset \mathfrak{M}_{\mathbb{Z}}(S)$ by taking non-negative linear combinations of all $\mu_{\mathcal{P}}$, where $\mathcal{P}$ runs over the set of all generalized polytopes in $\mathcal{P}(S)$.

Conjecture 24. The rational cone $\mathfrak{Pos}(S)$ is uniquely determined by the oriented matroid associated to S .

We conclude this section with the following question.

Problem 5. Is it possible to describe the extremal rays of $\mathfrak{Pos}(S)$?

One can easily show that a simplex from $\mathcal{P}(S)$ spans an extremal ray of $\mathfrak{Pos}(S)$ if and only if it does not contain any points of S distinct from its vertices. Problem 5 is apparently closely related to the problem of classification of combinatorial types of point arrangements, see e.g. [14] and references therein.

REFERENCES

- [1] T. V. Alekseevskaya, I. M. Gel'fand, and A. V. Zelevinskii, *Distribution of real hyperplanes and the partition function connected with it*, Dokl. Akad. Nauk SSSR **297** (1987), no. 6, 1289–1293. MR925528 (89m:05014)
- [2] T. Alekseyevskaya, *Bases in Systems of Simplices and Chambers*, ArXiv Mathematics e-prints (July 1997), available at [arXiv:math/9707218](https://arxiv.org/abs/math/9707218).
- [3] T. V. Alekseyevskaya, *Combinatorial bases in systems of simplices and chambers*, Proceedings of the 6th Conference on Formal Power Series and Algebraic Combinatorics (New Brunswick, NJ, 1994), 1996, pp. 15–37. MR1417284 (97k:52016)
- [4] M. Andersson, M. Passare, and R. Sigurdsson, *Complex convexity and analytic functionals*, Progress in Mathematics, vol. 225, Birkhäuser Verlag, Basel, 2004. MR2060426 (2005a:32011)
- [5] V. Baldoni, N. Berline, J. A. De Loera, M. Köppe, and M. Vergne, *How to integrate a polynomial over a simplex*, Math. Comp. **80** (2011), no. 273, 297–325. MR2728981
- [6] A. I. Barvinok, *Exponential integrals and sums over convex polyhedra*, Funktsional. Anal. i Prilozhen. **26** (1992), no. 2, 64–66. MR1173086 (93g:32052)
- [7] A. Barvinok, *Integer points in polyhedra*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2008. MR2455889 (2011a:52001)
- [8] M. Beck and S. Robins, *Computing the continuous discretely: integer-point enumeration in polyhedra*, Undergraduate Texts in Mathematics, Springer, New York, 2007. MR2271992 (2007h:11119)
- [9] P. Billingsley, *Convergence of probability measures*, Second, Wiley Series in Probability and Statistics: Probability and Statistics, John Wiley & Sons Inc., New York, 1999. A Wiley-Interscience Publication. MR1700749 (2000e:60008)
- [10] M. Bôcher, *Introduction to higher algebra*, Dover Publications Inc., New York, 1964. MR0172882 (30 #3098)
- [11] M. Brion, *Points entiers dans les polyèdres convexes*, Ann. Sci. École Norm. Sup. (4) **21** (1988), no. 4, 653–663. MR982338 (90d:52020)
- [12] M. A. Brodsky, *On the uniqueness of the inverse potential problem for homogeneous polyhedrons*, SIAM J. Appl. Math. **46** (1986), no. 2, 345–350. MR833481 (87k:86009b)
- [13] J. A. De Loera, J. Rambau, and F. Santos, *Triangulations*, Algorithms and Computation in Mathematics, vol. 25, Springer-Verlag, Berlin, 2010. Structures for algorithms and applications. MR2743368
- [14] K. Fukuda, H. Miyata, and S. Moriyama, *Complete enumeration of small realizable oriented matroids*, ArXiv e-prints (Apr. 2012), available at [1204.0645](https://arxiv.org/abs/1204.0645).

- [15] G. H. Golub, P. Milanfar, and J. Varah, *A stable numerical method for inverting shape from moments*, SIAM J. Sci. Comput. **21** (1999/00), no. 4, 1222–1243 (electronic). MR1740393 (2000m:65048)
- [16] N. Gravin, J. B. Lasserre, D. V. Pasechnik, and S. Robins, *The inverse moment problem for convex polytopes*, Vol. 48, 2011. <http://arxiv.org/abs/1106.5723>.
- [17] B. Gustafsson, C. He, P. Milanfar, and M. Putinar, *Reconstructing planar domains from their moments*, Inverse Problems **16** (2000), no. 4, 1053–1070. MR1776483 (2001k:44010)
- [18] J. Lawrence, *Polytope volume computation*, Math. Comp. **57** (1991), no. 195, 259–271. MR1079024 (91j:52019)
- [19] J. E. McCarthy and M. Putinar, *Positivity aspects of the Fantappiè transform*, J. Anal. Math. **97** (2005), 57–82. MR2274973 (2008m:46060)
- [20] P. Milanfar, M. Putinar, J. Varah, B. Gustafsson, and G. H. Golub, *Shape reconstruction from moments: theory, algorithms, and applications*, Society of photo-optical instrumentation engineers (spie) conference series, Nov. 2000, pp. 406–416.
- [21] I. Pak, *Lectures on discrete and polyhedral geometry*, UCLA, 2008. <http://www.math.ucla.edu/~pak/book.htm>.
- [22] D. Pasechnik and B. Shapiro, *On polygonal measures with vanishing harmonic moments*, ArXiv e-prints (Sep. 2012), available at 1209.4014.
- [23] A. V. Pukhlikov and A. G. Khovanskii, *The Riemann-Roch theorem for integrals and sums of quasipolynomials on virtual polytopes*, Algebra i Analiz **4** (1992), no. 4, 188–216. MR1190788 (94c:14044)
- [24] M. Putinar and C. Scheiderer, *Multivariate moment problems: geometry and indeterminateness*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **5** (2006), no. 2, 137–157. MR2244695 (2007e:44011)
- [25] W. Rudin, *Real and complex analysis*, Third Edition, McGraw-Hill Book Co., New York, 1987. MR924157 (88k:00002)
- [26] E. Schönhardt, *Über die Zerlegung von Dreieckspolyedern in Tetraeder*, Math. Ann. **98** (1928), no. 1, 309–312. MR1512408
- [27] V. N. Strakhov and M. A. Brodsky, *On the uniqueness of the inverse logarithmic potential problem*, SIAM J. Appl. Math. **46** (1986), no. 2, 324–344. MR833480 (87k:86009a)

SCHOOL OF PHYSICAL AND MATHEMATICAL SCIENCES, NANYANG TECHNOLOGICAL UNIVERSITY, 21 NANYANG LINK, 637371 SINGAPORE
E-mail address: ngravin@pmail.ntu.edu.sg

SCHOOL OF PHYSICAL AND MATHEMATICAL SCIENCES, NANYANG TECHNOLOGICAL UNIVERSITY, 21 NANYANG LINK, 637371 SINGAPORE
E-mail address: dima@ntu.edu.sg

DEPARTMENT OF MATHEMATICS, STOCKHOLM UNIVERSITY, SE-106 91 STOCKHOLM, SWEDEN
E-mail address: shapiro@math.su.se

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI 48824-1027, USA
E-mail address: mshapiro@math.msu.edu